

Symmetry solutions of the fifth-order non-linear equation with temporal and spatial dispersion terms

Zehra Pinar Izgi¹ and Subhadarshan Sahoo^{2,*}

¹Tekirdağ Namık Kemal University, Faculty of Arts and Science, Department of Mathematics, 59030 Merkez-Tekirdağ, Turkey.

²Department of Mathematics and Applied Statistics, School of Applied Sciences, Kalinga Institute of Industrial Technology, Deemed to be University, Bhubaneswar-751024, Odisha, India.

Abstract

The fifth-order non-linear partial differential equations (NPDEs) with temporal and spatial dispersion have been seen in the literature to model optical communication, fluid mechanics, condensed matter, electromagnetic phenomena, and the propagation of pulses in optical fibers. The considered model is not an easily solvable equation because of temporal and spatial dispersion. Our main aim is to obtain the explicit form of exact solutions via a combination of Lie group transformation and ansatz-based methods. Finally, the obtained results, which are novel solutions in explicit form, have been presented to illustrate the physical nature by means of three-dimensional plots, with a discussion of the results.

Keywords. Fifth-order non-linear equation, Exact solutions, Lie group transformation, Ansatz-based methods, Solitons.

2010 Mathematics Subject Classification. 35CXX, 37K40, 35QXX..

1. INTRODUCTION

Mathematical models are arising in engineering and applied science as non-linear partial differential equations (NPDEs) [7, 11, 20–22, 24]. The exact solutions of them play a major role in explaining the models and whether satisfy the process. Most of the described models are integrable, for which the exact solutions can be mentioned in various literature [6, 11–14, 18, 20, 22, 25].

Such models include the Korteweg–de Vries (KdV) equation, the nonlocal modified KdV equation, the nonlinear derivative Schrödinger equation, the Kadomtsev–Petviashvili (KP) equation, and the Camassa–Holm (CH) equation, and they have a variety of application areas [3, 7, 10, 11, 15, 17, 19–22, 25]. Lately, in the literature, fifth-order NPDEs include the temporal and spatial distribution terms applied to model optical communication, fluid mechanics, condensed matter, and the propagation of pulses in optical fibers. For this reason, we have considered here the fifth-order NPDE [21] as follows:

$$u_{ttt} - \lambda_1 u_{txxxx} - \lambda_2 u_{xxt} - \lambda_3 (u_x u_t)_{xx} - \lambda_4 (u_x u_{tx})_x = 0, \quad (1.1)$$

where $\lambda_i \neq 0$ ($i = 1, 2, 3, 4$) are parameters, u_{txxxx} is the spatial derivative and u_{ttt} is third-order dispersion causing a new form of impact instability [24]. Eq. (1.1) was considered for $\lambda_2 = 0$ and $\lambda_3 = \lambda_4 = 4$ by Wazwaz [21, 23], where Hirota's method is considered for solving the nonlinearity; additionally Wang et al. [20] considered Lie symmetry method. Besides Eq. (1.1) was considered for $\lambda_1 = 1, \lambda_2 = 1, \lambda_3 = \lambda_4 = 4$ and additionally six fifth-order non-linear equations were considered by Wazwaz [22].

Moreover, for the general case, the number of fifth-order equations is limitless. It is known that Eq. (1.1) is not an easily solvable equation because of temporal and spatial dispersion, but it has a variety of travelling waves such as solitons, peakons, kinks, etc. [22]. The effect of the dispersion term on the behavior is that it has a kink solution that

Received: 04 December 2024; Accepted: 20 July 2025.

* Corresponding author. Email: subha.bapi25@gmail.com.

changes it from one asymptotic state to another. So, the soliton structure of Eq.(1.1) is more complicated because of the spatial and dispersion terms.

The general case of the fifth-order non-linear equation Eq. (1.1) is considered in this work. As mentioned, due to the difficulty of both nonlinearity and dispersion, classical methods known in the literature are inadequate.

Our main contribution is to give the exact solution of the most general form of the equation Eq. (1.1) by a combination of the Lie group transformation and the ansatz-based method. To the best of our knowledge, these solutions are the most general solutions; they are presented here.

2. LIE TRANSFORMATIONS

One of the well-known and useful tools for reducing non-linear differential equations is the Lie transformation method [1–5, 8, 9, 15–17]. Briefly, with the help of the Lie transformations, which are invariant transformations, the considered linear or non-linear differential equation can be reduced to a solvable equation or a lower-order equation.

A one-parameter group transformation is considered as:

$$\tilde{x}^i = f^i(x, \varepsilon), \quad \varepsilon \text{ is a canonical parameter,} \quad i = 1, 2, \dots, n. \quad (2.1)$$

When the Taylor's series of $f^i(x, \varepsilon)$ at $\varepsilon = 0$, where $O(\varepsilon)$ is neglected, is considered, the infinitesimal transformation holds as follows:

$$\tilde{x}^i \approx x^i + \varepsilon \xi^i(x), \quad x^i = f^i(x, 0), \quad \xi^i(x) = \left. \frac{\partial f^i(x, \varepsilon)}{\partial \varepsilon} \right|_{\varepsilon=0}, \quad i = 1, 2, \dots, n. \quad (2.2)$$

Geometrically, ξ^i determines the tangent field vector of the group infinitesimal generator of group generator, which is given as:

$$X_j = \sum_{i=1}^n \xi^i(x) \frac{\partial}{\partial x^i}, \quad j = 1, \dots, n. \quad (2.3)$$

The generalized infinitesimal transformation is $X = \sum_{j=1}^n a_j X_j$, where $a_j (j = 1, \dots, n)$ are parameters, each considered sum satisfies the group properties.

3. METHODOLOGY

Sometimes the reduced equation is not generally solvable analytically with the theory of differential equations. So, in this work, modifications of the well-known AEM [10–14] are considered to obtain the exact solutions. Moreover, in the next section, the application of the methodology is presented in detail.

3.1. Solutions and Explanations: In this section, firstly the Lie transformations of the generalized fifth-order NPDE Eq. (1.1) are obtained, and then, according to the reduced equation, it is solved directly via the classical theory of differential equations or modifications of the well-known AEM.

Applying the given procedure, the determining equations have one solution set where $\Phi(x)$ is an arbitrary function

$$\zeta_x = C_2, \quad \zeta_t = C_1, \quad \zeta_u = \Phi(x). \quad (3.1)$$

As seen from the solution set, Lie group transformations change according to an arbitrary $F(x)$ function. Determination of $\Phi(x)$ is an open research area for physicists.

Depending on Eq.(3.1), two single-parameter groups are obtained:

$$\begin{aligned} &\text{For } C_1 = 1, C_2 = 0, \\ &X_1 = \frac{\partial}{\partial t} + \Phi(x) \frac{\partial}{\partial u}, \end{aligned} \quad (3.2)$$



and

$$\text{For } C_1 = 0, C_2 = 1, \\ X_2 = \frac{\partial}{\partial x} + \Phi(x) \frac{\partial}{\partial u}. \quad (3.3)$$

Case 1. Considering $X_2 = \frac{\partial}{\partial x} + \frac{1}{x} \frac{\partial}{\partial u}$ for $\Phi(x) = \frac{1}{x}$, the general form of solution $u(x, t) = \Theta(t)x$ is obtained. When the general solution is substituted in Eq.(1.1) with the $\lambda_3 = \lambda_4 = 4, \lambda_2 = 0, \lambda_1 = 1, \Theta(t) = \frac{c_1}{2}t^2 + c_2t + c_3$ is obtained. Hence, $u(x, t) = (\frac{c_1}{2}t^2 + c_2t + c_3)x$ is the generalized solution.

Case 2. The generalized infinitesimal transformation is $X = a_1X_1 + a_2X_2$. For this case, $a_1 = a_2 = 1$ i.e. $X = X_1 + X_2$ for $\Phi(x) = \frac{1}{x}$ is considered for the generalized type of Eq.(1.1) with the $\lambda_3 = \lambda_4 \neq 0, \lambda_1 \neq 0, \lambda_2 \neq 0$. Therefore, the solution is $u(x, t) = xH(\zeta), \zeta = -x + t$ and the reduced equation is held as

$$-\lambda_1 x H^{(5)} + 4\lambda_1 H^{(4)} + 6\lambda_3 (H')^2 + (\lambda_2(2-x) - 16x\lambda_3 H' + 4\lambda_3 H) H'' \\ + x(1 - \lambda_2 + \lambda_3(3xH' - 2H)) H''' = 0. \quad (3.4)$$

Firstly, to solve Eq.(3.4), the method of reducing order, which depends on the theory of differential equations, is considered. Therefore, the coefficient of the third and second derivatives of $H(\zeta)$ is assumed to be zero. As a result,

$$H(\zeta) = -\frac{(-2 + \lambda_2 x)\zeta}{10\lambda_3 x} - \frac{-16x + 10\lambda_2 x + 3\lambda_2 x^2 + 4\zeta - 2\zeta\lambda_2 x}{20\lambda_3 x}, \quad (3.5)$$

is obtained. The solution of Eq.(1.1) is

$$u(x, t) = x \left(-\frac{(-2 + \lambda_2 x)(-x + t)}{10\lambda_3 x} - \frac{-16x + 10\lambda_2 x + 3\lambda_2 x^2 + 4(-x + t) - 2(-x + t)\lambda_2 x}{20\lambda_3 x} \right), \quad (3.6)$$

where λ_3 and λ_2 are parameters which have physical meanings.

Case 3. The generalized infinitesimal transformation is $X = a_1X_1 + a_2X_2$. For this case, $a_1 = -1, a_2 = 1$ i.e. $X = X_1 + X_2$ for $\Phi(x) = \frac{1}{x}$ is considered for the generalized type of Eq.(1.1) with the $\lambda_3 \neq \lambda_4, \lambda_1, \lambda_2$ and all are different from zero. Therefore, $u(x, t) = xH(\zeta), \zeta = x + t$ is considered and the reduced equation is held as

$$-\lambda_1 x H^{(5)} - 4\lambda_1 H^{(4)} + x(1 - \lambda_2 - x(2\lambda_3 + \lambda_2)) H' - (\lambda_3 + \lambda_4) H H'' \\ + (-2\lambda_2 - 2x^2\lambda_3 - x(11\lambda_3 + 5\lambda_4)) H' - (2\lambda_3 + 2\lambda_4) H - (4\lambda_3 + \lambda_4(x^2 + 2)) H'' H'' = 0. \quad (3.7)$$

For the simplicity of Eq.(3.7), $\lambda_4 = -2\lambda_3$ is considered. To solve Eq.(3.7), the Hermite approximation method is considered. The first step is to determine the ansatz via the balancing principle. Hence, the ansatz is $H(\zeta) = \sum_{i=0}^N a_i z(\zeta)^i$, where N is determined via the balancing principle, a_i are the parameters that will be determined by solving the algebraic system, and $z(\zeta)$ is the solution of the Hermite differential (HD) equation $z'' - 2\zeta z' - \lambda z = 0$, which is the auxiliary equation. For the second step, the ansatz and the HD equation are substituted into Eq.(3.7), and a system of equations is obtained by polynomial equivalence, and the parameters are its solutions.

For Eq.(3.7), $H(\zeta) = \sum_{i=0}^1 a_i z(\zeta)^i$ is determined by using balancing principle. Applying the given procedure, the solution sets satisfying our goals for the parameters are proposed in Table 1, and others give trivial or meaningless solutions.

For the first set, the solution is

$$u(x, t) = \frac{2(x+t)x^3\lambda_2 - 8\lambda_1(x+t)^2 - 6\lambda_1(x+t)x - 4\lambda_1 - 4\lambda_1(x+t)^3x + x(x+t) - x\lambda_2(x+t) - \lambda_2}{(x+t)(x(x+t)+1)\lambda_2} \\ + \frac{(x(x+t)+1)\left(c_1(x+t)KummerM\left(0, \frac{3}{2}, (x+t)^2\right) + c_2(x+t)KummerU\left(0, \frac{3}{2}, (x+t)^2\right)\right)}{2x}. \quad (3.8)$$



TABLE 1. The solution sets of parameters for Case 3.

Set 1	Set 2
$\lambda = -2,$ $a_1 = \frac{x\zeta+1}{2x^2},$ $\lambda_3 = -2x\lambda_2\zeta,$ $a_0 = \frac{2\zeta x^3\lambda_2 - 8\lambda_1\zeta^2 - 6\lambda_1\zeta x - 4\lambda_1 - 4\lambda_1\zeta^3 x + x\zeta - x\lambda_2\zeta - \lambda_2}{\zeta(x\zeta+1)\lambda_2 x}$	$\lambda = \frac{2x^2-3x\zeta+1}{x^2}, a_1 = \frac{x\zeta+1}{2x^2-3x\zeta+1},$ $\lambda_2 = \frac{2(x^2\lambda_1\zeta(2x-\zeta) - \lambda_1(6\zeta x+1)+x)}{x^4(x\zeta+1)},$ $\lambda_3 = \frac{x^2\lambda_1\zeta(2x-\zeta) - \lambda_1(6\zeta x+1)+x}{x^4},$ $a_0 = -\frac{4x^5\lambda_1\zeta(\zeta^2+1) - x^4(\zeta x+1) + x^3\lambda_1(14\zeta+4x+15x\zeta^2)}{(x\zeta+1)(x^2\lambda_1\zeta(2x-\zeta) - \lambda_1(6\zeta x+1)+x^2)} + \frac{-x^2\lambda_1(1+2\zeta^2) - 2\lambda_1(6\zeta x+1)+2x^2}{(x\zeta+1)(x^2\lambda_1\zeta(2x-\zeta) - \lambda_1(6\zeta x+1)+x^2)}$

Case 4. The generalized infinitesimal transformation is $X = X_1 + X_2$ and $\Phi(x) = 0$ is considered for the generalized type of Eq.(1.1) with the $\lambda_3 = \lambda_4 \neq 0, \lambda_1 \neq 0, \lambda_2 \neq 0$. Therefore, the solution is $u(x, t) = H(\zeta), \zeta = x + t$ and the reduced equation is held as

$$-\lambda_1 H^{(5)} - 3\lambda_3 (H'')^2 + (1 - \lambda_2 - 3\lambda_3 H') H''' = 0. \quad (3.9)$$

The given procedure of the Hermite approximation method is applied, and as a result, the parameters are obtained

$$a_1 = \frac{-4\lambda_1(4\zeta^2 - 1)}{3\lambda_3((c_1\kappa_1 + c_2\kappa_2)(7 + 10\zeta^2) - c_1\kappa_3(9 + 12\zeta^2) + c_2\kappa_4)}, \quad \lambda_2 = -4\lambda_1\zeta^2 - 2\lambda_1 + 1, \quad (3.10)$$

where $\kappa_1 = KummerM(-2\zeta^2, \frac{3}{2}, \zeta^2)$, $\kappa_2 = KummerU(-2\zeta^2, \frac{3}{2}, \zeta^2)$, $\kappa_3 = KummerM(1 - 2\zeta^2, \frac{3}{2}, \zeta^2)$ and $\kappa_4 = KummerU(1 - 2\zeta^2, \frac{3}{2}, \zeta^2)$.

Substituting the parameters and $z(\zeta)$ into the ansatz, the general solution is obtained.

4. RESULT AND DISCUSSION

The dynamics of wave propagation in the pertinent media have been covered in this section. The aforementioned solutions were thoroughly investigated, discussed, and shown using three-dimensional graphs and numerical values applied to the unknown parameters.

4.1. Graphical representation of the solutions. Here, the three dimensional graphical representation of the plot for Case 3 and Case 4 is given by Figures 1 and 2 as follows:

4.2. Dynamics of the solutions. This section explains the physical meaning of the several solution graphs that were successfully achieved using numerical simulation of the unknown components. Moreover, the dynamic characteristics of the distinct waveforms that were acquired and their application to the pertinent media are covered in this section. The graphical picturization of Case 3 and Case 4 under distinct parametric conditions has been shown in Figures 1 and 2, showing the soliton that propagated throughout the medium with assigned different values of unknown parameters.

In the above section, the graphical analysis illustrates how the waves proceed without any information loss at each point in time, and these wave patterns arise from assigning discrete numerical values to the unknowns. Furthermore, it is evident from the description above how these waves are applicable in diverse environments and how they will act in a medium with distinct intensity backgrounds. Furthermore, these solutions retain their shapes during propagation in the medium, resulting in their durability and stability.

5. CONCLUSION

In this paper, the exact and explicit solutions of Eq.(1.1) via Lie symmetry analysis and in combination with a modification of the well-known AEM, where the Hermite approximation method is used, are obtained. The determination of the Lie group generators has an arbitrary function $F(x)$ that may have physical meaning. The considered model is not an easily solvable equation because of temporal and spatial dispersion. Our main contribution is to give the exact solution of the most general form of Eq.(1.1) by a combination of the Lie group transformation and the ansatz-based






FIGURE 1. The solution of Case 3 considering the parameters of Set 1 for $c_2 = 1$, $c_1 = 1$, $\lambda_2 = -3$, $\lambda_1 = -1$.




FIGURE 2. The solution of Case 4 for $a_0 = 1$, $c_1 = 1$, $c_2 = -1$, $\lambda = 1$.

method, supported by Figures. Finally, the newly obtained results have been presented in the explicit form with physical interpretation in three-dimensional plots to show the nature of the solutions. The scientific contribution of this work is related to the research area in solitary waves theory, where a new generalized fifth-order integrable equation is introduced and investigated.

FUNDING

No funding was received for the preparation of this manuscript.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

DATA AVAILABILITY

Data sharing is not applicable to this article, as no datasets were generated or analyzed during the current study.

ACKNOWLEDGEMENTS

The authors would also like to thank the anonymous referees and editor for all the valuable suggestions and comments which improved both the quality and clarity of the paper.

REFERENCES

- [1] W. F. Ames and W. N. Nucci *Analysis of fluid equations by group methods*, Journal of Engineering Mechanics, 20 (1985), 181-187.
- [2] G. W. Bluman and S. C. Anco *Symmetry and Integration Methods for Differential Equations*, Applied Mathematical Sciences, 154 (2002).
- [3] G. Guirao, H. M. Baskonus A. Kumar, M. S. Rawat, and G. Yel, *Complex Patterns to the (3+1)-Dimensional B-type Kadomtsev-Petviashvili-Boussinesq Equation*, Symmetry, 83 (2020), 12-17.



- [4] N. H. Ibragimov, *CRC Handbook of Lie Group Analysis of Differential Equations, Vol. 1: Symmetries, Exact Solutions and Conservation Laws*, CRC, Boca Raton, Florida, 1994.
- [5] N. H. Ibragimov, *Transformation Groups and Lie Algebras*, World Scientific Publishing, 2013.
- [6] H. F. Ismael, H. Bulut, and H. M. Baskonus, *Optical soliton solutions to the Fokas–Lenells equation via sine-Gordon expansion method and [Math Processing Error]-expansion method*, *Pramana - J Phys*, *94* (2020).
- [7] C. K. Kuo, *Resonant multi-soliton solutions to two fifth-order KdV equations via the simplified linear superposition principle*, *Modern Physics Letters B*, *33* (2019), 1950299.
- [8] P. J. Oliver *Applications of Lie Groups to differential Equations*, Springer Verlag, New York, 1993 .
- [9] T. Ozis and I. Aslan *Similarity solutions to Burgers' equation in terms of special functions of mathematical physics*, *Acta Physica Polonica B*, *48*(7) (2017), 1349-1369.
- [10] Z. Pinar and T. Oziş *Classical symmetry analysis and exact solutions for generalized Korteweg–de Vries models with variable coefficients*, *Int. J. Nonlin. Mech.*, *105* (2018), 99-104.
- [11] Z. Pinar and H. Kocak *Exact solutions for the third-order dispersive-Fisher equations*, *Nonlinear Dyn*, *91* (2018), 421-426.
- [12] Z. Pinar, *The Symmetry Analysis of Electrostatic Micro-Electromechanical System (MEMS)*, *Modern Physics Letters B*, *34* (2020), 2050199.
- [13] Z. Pinar, *The Combination of Conservation Laws and Auxiliary Equation Method*, *Int. J. Appl. Comput. Math*, *6* (2020).
- [14] Z. Pinar, *Simulations of surface corrugations of graphene sheets through the generalized graphene thermophoretic motion equation*, *International Journal of Computational Materials Science and Engineering*, *2050005* (2020).
- [15] S. Sahoo and S. S. Ray, *Analysis of Lie symmetries with conservation laws for the (3+1) dimensional time-fractional mKdV-ZK equation in ion-acoustic waves*, *Nonlinear Dyn*, *90* (2017), 1105-1113.
- [16] S. Sahoo and S. S. Ray, *Invariant analysis with conservation laws for the time fractional Drinfeld-Sokolov-Satsuma-Hirota equations by Lie group analysis*, *Chaos, Solitons & Fractals*, *104* (2017), 725-733.
- [17] S. Sahoo and S. S. Ray, *The conservation laws with Lie symmetry analysis for time fractional integrable coupled KdV-mKdV system*, *International Journal of Non-Linear Mechanics*, *98* (2017), 114-121.
- [18] Y. G. Sánchez, Z. Sabir, H. Günerhan, and H. M. Baskonus, *Analytical and Approximate Solutions of a Novel Nervous Stomach Mathematical Model*, *Discrete Dynamics in Nature and Society*, *2020* (2020), 5063271.
- [19] S. Singh, K. Sakkaravarthi, and K. Murugesan, *Lump and soliton on certain spatially-varying backgrounds for an integrable (3+1) dimensional fifth-order nonlinear oceanic wave model*, *Chaos, Solitons & Fractals*, *167* (2023), 113058.
- [20] G. W. Wang, X. G. Liu, and Y. Y. Zhang, *Symmetry reduction, exact solutions and conservation laws of a new fifth-order nonlinear integrable equation*, *Commun Nonlinear Sci Numer Simulat*, *18* (2013), 2313-2320.
- [21] A. M. Wazwaz, *A new fifth-order nonlinear integrable equation: multiple soliton solutions*, *Physica Scripta*, *83* (2011), 015012.
- [22] A. M. Wazwaz, *Kink solutions for three new fifth order nonlinear equations*, *Applied Mathematical Modelling*, *38* (2014), 110-118.
- [23] A. M. Wazwaz, *Multiple soliton solutions and other scientific solutions for a new Painlevé integrable fifth-order equation*, *Chaos, Solitons & Fractals*, *196* (2025), 116307.
- [24] G. Q. Xu and A. M. Wazwaz, *Bidirectional solitons and interaction solutions for a new integrable fifth-order nonlinear equation with temporal and spatial dispersion*, *Nonlinear Dynamics*, *101* (2020), 581-595.
- [25] G. Yel, H. M. Baskonus, and W. Gao, *New dark-bright soliton in the shallow water wave model*, *AIMS Mathematics*, *5*(4) (2020), 4027-4044.

