



## New midpoint type inequalities for generalized fractional integral

Hüseyin Budak\*, Hasan Kara, and Rabia Kapucu

Department of Mathematics, Faculty of Science and Arts, Düzce University, Düzce, Turkey.

### Abstract

In this paper, we first establish two new identities for differentiable function involving generalized fractional integrals. Then, by utilizing these equalities, we obtain some midpoint type inequalities involving generalized fractional integrals for mappings whose derivatives in absolute values are convex. We also give several results as special cases of our main results.

**Keywords.** Hermite-Hadamard inequality, Generalized fractional integral, Convex function.

**2010 Mathematics Subject Classification.** 26D15, 26B25, 26D10.

### 1. INTRODUCTION

In recent years, the Hermite-Hadamard inequality, which is the first fundamental result for convex mappings with a natural geometrical interpretation and many applications, has drawn attention much interest in elementary mathematics.

The inequalities discovered by C. Hermite and J. Hadamard for convex functions are considerable significance in the literature (see, e.g., [13, 29]). These inequalities state that if  $f : I \rightarrow \mathbb{R}$  is a convex function on the interval  $I$  of real numbers and  $\kappa_1, \kappa_2 \in I$  with  $\kappa_1 < \kappa_2$ , then

$$f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \leq \frac{1}{\kappa_2 - \kappa_1} \int_{\kappa_1}^{\kappa_2} f(\kappa) dx \leq \frac{f(\kappa_1) + f(\kappa_2)}{2}. \quad (1.1)$$

Both inequalities hold in the reversed direction if  $f$  is concave.

Over the last twenty years, the numerous studies have focused on to obtain new bound for left hand side and right side of the inequality (1.1). For some examples, please refer to ([6, 8, 11, 13, 14, 24, 31–34, 40]).

In the following we will give some necessary definitions and mathematical preliminaries of fractional calculus theory which are used further in this paper.

**Definition 1.1.** [23] Let  $f \in L_1[\kappa_1, \kappa_2]$ . The Riemann-Liouville integrals  $J_{\kappa_1+}^\alpha f$  and  $J_{\kappa_2-}^\alpha f$  of order  $\alpha > 0$  with  $\kappa_1 \geq 0$  are defined by

$$J_{\kappa_1+}^\alpha f(\kappa) = \frac{1}{\Gamma(\alpha)} \int_{\kappa_1}^{\kappa} (\kappa - \xi)^{\alpha-1} f(\xi) d\xi, \quad \kappa > \kappa_1$$

and

$$J_{\kappa_2-}^\alpha f(\kappa) = \frac{1}{\Gamma(\alpha)} \int_{\kappa}^{\kappa_2} (\xi - \kappa)^{\alpha-1} f(\xi) d\xi, \quad \kappa < \kappa_2$$

respectively. Here,  $\Gamma(\alpha)$  is the Gamma function and  $J_{\kappa_1+}^0 f(\kappa) = J_{\kappa_2-}^0 f(\kappa) = f(\kappa)$ .

Received: 17 July 2020 ; Accepted: 12 December 2020.

\* Corresponding author. Email: hsyn.budak@gmail.com.

It is remarkable that Sarikaya et al.[37] first give the following interesting integral inequalities of Hermite-Hadamard type involving Riemann-Liouville fractional integrals.

**Theorem 1.2.** *Let  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a positive function with  $0 \leq \kappa_1 < \kappa_2$  and  $f \in L_1[\kappa_1, \kappa_2]$ . If  $f$  is a convex function on  $[\kappa_1, \kappa_2]$ , then the following inequalities for fractional integrals hold:*

$$\begin{aligned} f\left(\frac{\kappa_1 + \kappa_2}{2}\right) &\leq \frac{\Gamma(\alpha + 1)}{2(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\kappa_1^+}^\alpha f(\kappa_2) + J_{\kappa_2^-}^\alpha f(\kappa_1) \right] \\ &\leq \frac{f(\kappa_1) + f(\kappa_2)}{2} \end{aligned} \quad (1.2)$$

with  $\alpha > 0$ .

Sarikaya and Yildirim also give the following Hermite-Hadamard type inequality for the Riemann-Liouville fractional integrals in [36].

**Theorem 1.3.** *Let  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a positive function with  $\kappa_1 < \kappa_2$  and  $f \in L_1[\kappa_1, \kappa_2]$ . If  $f$  is a convex function on  $[\kappa_1, \kappa_2]$ , then the following inequalities for fractional integrals hold:*

$$\begin{aligned} f\left(\frac{\kappa_1 + \kappa_2}{2}\right) &\leq \frac{2^{\alpha-1}\Gamma(\alpha + 1)}{(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^+}^\alpha f(\kappa_2) + J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^-}^\alpha f(\kappa_1) \right] \\ &\leq \frac{f(\kappa_1) + f(\kappa_2)}{2}. \end{aligned} \quad (1.3)$$

Whereupon Sarikaya et al. obtain the Hermite-Hadamard inequality for Riemann-Liouville fractional integrals, many authors have studied to generalize this inequality and establish Hermite-Hadamard inequality other fractional integrals such as  $k$ -fractional integral, Hadamard fractional integrals, Katugampola fractional integrals, Conformable fractional integrals, etc. For some of them, please see ([1-3, 7, 9, 12], [15]-[21], [25, 26, 28, 30, 38, 39], [41]-[46]).

## 2. NEW GENERALIZED FRACTIONAL INTEGRAL OPERATORS

In this section we summarize the generalized fractional integrals defined by Sarikaya and Ertuğral in [35]. Let's define a function  $\varphi : [0, \infty) \rightarrow [0, \infty)$  satisfying the following conditions :

$$\int_0^1 \frac{\varphi(\xi)}{\xi} d\xi < \infty.$$

We define the following left-sided and right-sided generalized fractional integral operators, respectively, as follows:

$${}_{\kappa_1^+} I_\varphi f(\kappa) = \int_{\kappa_1}^{\kappa} \frac{\varphi(\kappa - \xi)}{\kappa - \xi} f(\xi) d\xi, \quad \kappa > \kappa_1, \quad (2.1)$$

$${}_{\kappa_2^-} I_\varphi f(\kappa) = \int_{\kappa}^{\kappa_2} \frac{\varphi(\xi - \kappa)}{\xi - \kappa} f(\xi) d\xi, \quad \kappa < \kappa_2. \quad (2.2)$$

The most important feature of generalized fractional integrals is that they generalize some types of fractional integrals such as Riemann-Liouville fractional integrals,  $k$ -Riemann-Liouville fractional integrals, Katugampola fractional integrals, conformable fractional integrals, Hadamard fractional integrals, etc. These important special cases of the integral operators (2.1) and (2.2) are mentioned below.

i) If we take  $\varphi(\xi) = \xi$ , the operator (2.1) and (2.2) reduce to the Riemann integral as follows:

$$\begin{aligned} I_{\kappa_1^+} f(\kappa) &= \int_{\kappa_1}^{\kappa} f(\xi) d\xi, \quad \kappa > \kappa_1, \\ I_{\kappa_2^-} f(\kappa) &= \int_{\kappa}^{\kappa_2} f(\xi) d\xi, \quad \kappa < \kappa_2. \end{aligned}$$



ii) If we take  $\varphi(\xi) = \frac{\xi^\alpha}{\Gamma(\alpha)}$ , the operator (2.1) and (2.2) reduce to the Riemann-Liouville fractional integral as follows:

$$I_{\kappa_1^+}^\alpha f(\kappa) = \frac{1}{\Gamma(\alpha)} \int_{\kappa_1}^\kappa (\kappa - \xi)^{\alpha-1} f(\xi) d\xi, \quad \kappa > \kappa_1,$$

$$I_{\kappa_2^-}^\alpha f(\kappa) = \frac{1}{\Gamma(\alpha)} \int_\kappa^{\kappa_2} (\xi - \kappa)^{\alpha-1} f(\xi) d\xi, \quad \kappa < \kappa_2.$$

iii) If we take  $\varphi(\xi) = \frac{1}{k\Gamma_k(\alpha)} \xi^{\frac{\alpha}{k}}$ , the operator (2.1) and (2.2) reduce to the  $k$ -Riemann-Liouville fractional integral as follows:

$$I_{\kappa_1^+,k}^\alpha f(\kappa) = \frac{1}{k\Gamma_k(\alpha)} \int_{\kappa_1}^\kappa (\kappa - \xi)^{\frac{\alpha}{k}-1} f(\xi) d\xi, \quad \kappa > \kappa_1,$$

$$I_{\kappa_2^-,k}^\alpha f(\kappa) = \frac{1}{k\Gamma_k(\alpha)} \int_\kappa^{\kappa_2} (\xi - \kappa)^{\frac{\alpha}{k}-1} f(\xi) d\xi, \quad \kappa < \kappa_2$$

where

$$\Gamma_k(\alpha) = \int_0^\infty \xi^{\alpha-1} e^{-\frac{\xi^k}{k}} d\xi, \quad \mathcal{R}(\alpha) > 0$$

and

$$\Gamma_k(\alpha) = k^{\frac{\alpha}{k}-1} \Gamma\left(\frac{\alpha}{k}\right), \quad \mathcal{R}(\alpha) > 0; k > 0$$

are given by Mubeen and Habibullah in [27].

iv) If we take  $\varphi(\xi) = \xi(\kappa - \xi)^{\alpha-1}$ , the operator (2.1) reduces to the conformable fractional operators as follows:

$$I_{\kappa_1}^\alpha f(\kappa) = \int_{\kappa_1}^\kappa \xi^{\alpha-1} f(\xi) d\xi = \int_{\kappa_1}^\kappa f(\xi) d_\alpha \xi, \quad \kappa > \kappa_1, \alpha \in (0, 1)$$

is given by Khalil et al. in [22].

Sarikaya and Ertuğral also establish the following Hermite-Hadamard inequality for the generalized fractional integral operators:

**Theorem 2.1.** *Let  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a convex function on  $[\kappa_1, \kappa_2]$  with  $\kappa_1 < \kappa_2$ , then the following inequalities for fractional integral operators hold*

$$f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \leq \frac{1}{2\Psi(1)} [\kappa_{1+} I_\varphi f(\kappa_2) + \kappa_{2-} I_\varphi f(\kappa_1)] \leq \frac{f(\kappa_1) + f(\kappa_2)}{2} \tag{2.3}$$

where the mapping  $\Psi : [0, 1] \rightarrow \mathbb{R}$  is defined by

$$\Psi(\kappa) = \int_0^\kappa \frac{\varphi((\kappa_2 - \kappa_1)\xi)}{\xi} d\xi.$$

For more recent results related to generalized fractional integral inequalities see, ([4, 5, 10, 35]). In this paper, we obtain some midpoint type inequality for the generalized fractional integrals.

### 3. TWO NEW EQUALITIES FOR GENERALIZED FRACTIONAL INTEGRALS

Firstly, we give the following lemmas which will be used frequently in next section.



**Lemma 3.1.** *Let  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(\kappa_1, \kappa_2)$  with  $\kappa_1 < \kappa_2$ . If  $f' \in L[\kappa_1, \kappa_2]$ , then we have the following equality for generalized fractional integrals*

$$\begin{aligned} & \frac{1}{\Lambda(0) + \Delta(0)} \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \\ &= \frac{(\kappa - \kappa_1)}{\Lambda(0) + \Delta(0)} \int_0^1 \Lambda(\xi) f'(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \\ & \quad - \frac{(\kappa_2 - \kappa)}{\Lambda(0) + \Delta(0)} \int_0^1 \Delta(\xi) f'(\xi \kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \end{aligned} \quad (3.1)$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\Lambda$  and  $\Delta$  are defined by

$$\begin{aligned} \Lambda(\xi) &= \int_{\xi}^1 \frac{\varphi((\kappa - \kappa_1)u)}{u} du \\ \Delta(\xi) &= \int_{\xi}^1 \frac{\varphi((\kappa_2 - \kappa)u)}{u} du. \end{aligned} \quad (3.2)$$

*Proof.* Integrating by parts, we have

$$\begin{aligned} I_1 &= \int_0^1 \Lambda(\xi) f'(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \\ &= \left[ \Lambda(\xi) \frac{f(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))}{\kappa - \kappa_1} \right]_0^1 + \int_0^1 \frac{f(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))}{\kappa - \kappa_1} \frac{\varphi((\kappa - \kappa_1)\xi)}{\xi} d\xi. \end{aligned}$$

Using the change of variable  $t\kappa_2 + (1 - t)(\kappa_1 + \kappa_2 - \kappa) = \varsigma$ , we get,

$$\begin{aligned} I_1 &= -\Lambda(0) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa - \kappa_1} + \frac{1}{\kappa - \kappa_1} \int_{\kappa_1 + \kappa_2 - \kappa}^{\kappa_2} f(\varsigma) \frac{\varphi(\varsigma - (\kappa_1 + \kappa_2 - \kappa))}{\varsigma - (\kappa_1 + \kappa_2 - \kappa)} d\varsigma \\ &= -\Lambda(0) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa - \kappa_1} + \frac{1}{\kappa - \kappa_1} {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa). \end{aligned}$$

Similarly, we obtain

$$\begin{aligned} I_2 &= \int_0^1 \Delta(\xi) f'(\xi \kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \\ &= \Delta(0) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa_2 - \kappa} - \frac{1}{\kappa_2 - \kappa} {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa). \end{aligned}$$

If we multiply  $I_1$  and  $I_2$  by  $\frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)}$  and  $-\frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)}$ , respectively, then we get

$$\frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} I_1 - \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} I_2 = \frac{1}{\Lambda(0) + \Delta(0)} \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa)$$

which completes the proof.  $\square$



**Lemma 3.2.** *The assumptions of Lemma 3.1 hold. Then we have the following equality for generalized fractional integrals*

$$\begin{aligned} & \frac{1}{\Psi(1) + \Theta(1)} \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \\ &= \frac{(\kappa - \kappa_1)}{\Psi(1) + \Theta(1)} \int_0^1 \Psi(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2) d\xi \\ &+ \frac{(\kappa_2 - \kappa)}{\Psi(1) + \Theta(1)} \int_0^1 \Theta(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1) d\xi \end{aligned} \tag{3.3}$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\Psi$  and  $\Theta$  are defined by

$$\begin{aligned} \Psi(\xi) &= \int_0^\xi \frac{\varphi((\kappa - \kappa_1)u)}{u} du \\ \Theta(\xi) &= \int_0^\xi \frac{\varphi((\kappa_2 - \kappa)u)}{u} du. \end{aligned} \tag{3.4}$$

*Proof.* Integrating by parts, we have

$$\begin{aligned} I_3 &= \int_0^1 \Psi(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2) d\xi \\ &= \Psi(\xi) \frac{f(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2)}{\kappa_1 - \kappa} \Big|_0^1 - \int_0^1 \frac{f(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2)}{\kappa_1 - \kappa} \frac{\varphi((\kappa - \kappa_1)\xi)}{\xi} d\xi. \end{aligned}$$

Using change of variable  $\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2 = \tau$ , we get,

$$\begin{aligned} I_3 &= -\Psi(1) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa - \kappa_1} + \frac{1}{\kappa - \kappa_1} \int_{\kappa_1 + \kappa_2 - \kappa}^{\kappa_2} f(\tau) \frac{\varphi(\kappa_2 - \tau)}{\kappa_2 - \tau} d\tau \\ &= -\Psi(1) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa - \kappa_1} + \frac{1}{\kappa - \kappa_1} {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2). \end{aligned}$$

Similarly, we get

$$\begin{aligned} I_4 &= \int_0^1 \Theta(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1) d\xi \\ &= \Theta(1) \frac{f(\kappa_1 + \kappa_2 - \kappa)}{\kappa_2 - \kappa} - \frac{1}{\kappa_2 - \kappa} {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1). \end{aligned}$$

If we multiply  $I_3$  and  $I_4$  by  $\frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)}$  and  $-\frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)}$ , respectively, then we get,

$$\frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} I_3 - \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} I_4 = \frac{1}{\Psi(1) + \Theta(1)} \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa)$$

which completes the proof. □



## 4. MIDPOINT TYPE INEQUALITIES FOR GENERALIZED FRACTIONAL INTEGRALS

In this section, utilizing the identities obtained previous section, we establish some midpoint type inequalities for generalized fractional integrals.

**Theorem 4.1.**  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(\kappa_1, \kappa_2)$  with  $0 \leq \kappa_1 < \kappa_2$ . If  $|f'|$  is convex on  $[\kappa_1, \kappa_2]$ , then we have the following inequality for generalized fractional integrals

$$\left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \quad (4.1)$$

$$\begin{aligned} &\leq \frac{1}{\Lambda(0) + \Delta(0)} \left\{ (\kappa - \kappa_1) \left| f'(\kappa_2) \right| \int_0^1 \Lambda(\xi) \xi d\xi + (\kappa_2 - \kappa) \left| f'(\kappa_1) \right| \int_0^1 \Delta(\xi) \xi d\xi \right. \\ &\quad \left. + \left| f'(\kappa_1 + \kappa_2 - \kappa) \right| \int_0^1 (1 - \xi) [(\kappa - \kappa_1) \Lambda(\xi) + (\kappa_2 - \kappa) \Delta(\xi)] d\xi \right\} \quad (4.2) \end{aligned}$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\Lambda$  and  $\Delta$  are defined as in (3.2).

*Proof.* By the Lemma 3.1, we have

$$\begin{aligned} &\left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ &\leq \frac{(\kappa - \kappa_1)}{\Lambda(0) + \Delta(0)} \left| \int_0^1 \Lambda(\xi) f'(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \right| \\ &\quad + \frac{(\kappa_2 - \kappa)}{\Lambda(0) + \Delta(0)} \left| \int_0^1 \Delta(\xi) f'(\xi \kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa)) d\xi \right| \\ &\leq \frac{(\kappa - \kappa_1)}{\Lambda(0) + \Delta(0)} \int_0^1 |\Lambda(\xi) f'(\xi \kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))| d\xi \\ &\quad + \frac{(\kappa_2 - \kappa)}{\Lambda(0) + \Delta(0)} \int_0^1 |\Delta(\xi) f'(\xi \kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))| d\xi. \end{aligned}$$



Using convexity of  $|f'|$ , we obtain

$$\begin{aligned} & \left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{(\kappa - \kappa_1)}{\Lambda(0) + \Delta(0)} \left\{ \int_0^1 \Lambda(\xi) \xi |f'(\kappa_2)| + (1 - \xi) |f'(\kappa_1 + \kappa_2 - \kappa)| d\xi \right. \\ & \quad \left. + \frac{(\kappa_2 - \kappa)}{\Lambda(0) + \Delta(0)} \int_0^1 \Delta(\xi) \xi |f'(\kappa_1)| + (1 - \xi) |f'(\kappa_1 + \kappa_2 - \kappa)| d\xi \right\} \\ & = \frac{1}{\Lambda(0) + \Delta(0)} \times \left\{ (\kappa - \kappa_1) |f'(\kappa_2)| \int_0^1 \Lambda(\xi) \xi d\xi + (\kappa_2 - \kappa) |f'(\kappa_1)| \int_0^1 \Delta(\xi) \xi d\xi \right. \\ & \quad \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \int_0^1 (1 - \xi) [(\kappa - \kappa_1) \Lambda(\xi) + (\kappa_2 - \kappa) \Delta(\xi)] d\xi \right\}. \end{aligned}$$

This completes the proof. □

**Corollary 4.2.** Under assumption of Theorem 4.1 with  $\varphi(\xi) = \xi$ , we have the following inequality

$$\begin{aligned} \left| \frac{1}{\kappa_2 - \kappa_1} \int_{\kappa_1}^{\kappa_2} f(\xi) d\xi - f(\kappa_1 + \kappa_2 - \kappa) \right| & \leq \frac{1}{\kappa_2 - \kappa_1} \left\{ \frac{1}{6} \left[ (\kappa - \kappa_1)^2 |f'(\kappa_2)| + (\kappa_2 - \kappa)^2 |f'(\kappa_1)| \right] \right. \\ & \quad \left. + \frac{1}{3} |f'(\kappa_1 + \kappa_2 - \kappa)| [(\kappa - \kappa_1)^2 + (\kappa_2 - \kappa)^2] \right\}. \end{aligned} \tag{4.3}$$

**Remark 4.3.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.2, then we have the following inequality

$$\left| \frac{1}{\kappa_2 - \kappa_1} \int_{\kappa_1}^{\kappa_2} f(\xi) d\xi - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{\kappa_2 - \kappa_1}{8} \left[ |f'(\kappa_1)| + |f'(\kappa_2)| \right]$$

which is proved by Kirmaci in [24].

**Corollary 4.4.** Under assumption of Theorem 4.1 with  $\varphi(\xi) = \frac{\xi^\alpha}{\Gamma(\alpha)}$ , we have the following inequality

$$\begin{aligned} & \left| \frac{\Gamma(\alpha + 1)}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \times \left[ J_{\kappa_2^-}^\alpha f(\kappa_1 + \kappa_2 - \kappa) + J_{\kappa_1^+}^\alpha f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \frac{\alpha}{2(\alpha + 2)} \left\{ (\kappa - \kappa_1)^{\alpha+1} |f'(\kappa_2)| + (\kappa_2 - \kappa)^{\alpha+1} |f'(\kappa_1)| \right. \\ & \quad \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \frac{\alpha + 3}{\alpha + 1} [(\kappa_2 - \kappa)^{\alpha+1} + (\kappa - \kappa_1)^{\alpha+1}] \right\} \end{aligned} \tag{4.4}$$

**Remark 4.5.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.4, then we have the following inequality

$$\left| \frac{2^{\alpha-1} \Gamma(\alpha + 1)}{(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\kappa_2^-}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) + J_{\kappa_1^+}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{(\kappa_2 - \kappa_1)^\alpha}{4(\alpha + 1)} \left[ |f'(\kappa_1)| + |f'(\kappa_2)| \right]$$

which is given by Budak and Agarwal in [9, Corollary 2 for  $q = 1$ ].



**Corollary 4.6.** Under assumption of Theorem 4.1 with  $\varphi(\xi) = \frac{\xi^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ , we have the following inequality

$$\begin{aligned} & \left| \frac{\Gamma_k(\alpha + k)}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \times \left[ J_{\kappa_2^-, k}^{\alpha} f(\kappa_1 + \kappa_2 - \kappa) + J_{\kappa_1^+, k}^{\alpha} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^{\alpha/k} + (\kappa - \kappa_1)^{\alpha/k}} \\ & \times \left\{ (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} \frac{\alpha}{2(\alpha + 2k)} |f'(\kappa_2)| + (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} \frac{\alpha}{2(\alpha + 2k)} |f'(\kappa_1)| \right. \\ & \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \frac{\alpha(\alpha + 3k)}{2(\alpha + k)(\alpha + 2k)} \left[ (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} + (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} \right] \right\}. \end{aligned}$$

**Remark 4.7.** Under assumption of Corollary 4.6 with  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$ , we have the following inequality

$$\left| \frac{2^{\frac{\alpha}{k}-1} \Gamma_k(\alpha + k)}{(\kappa_2 - \kappa_1)^{\frac{\alpha}{k}}} \left[ J_{\kappa_2^-, k}^{\alpha} f\left(\frac{\kappa_1 + \kappa_2}{2}\right) + J_{\kappa_1^+, k}^{\alpha} f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{(\kappa_2 - \kappa_1)\alpha}{4(\alpha + k)} [|f'(\kappa_1)| + |f'(\kappa_2)|].$$

**Theorem 4.8.**  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(\kappa_1, \kappa_2)$  with  $0 \leq \kappa_1 < \kappa_2$ . If  $|f'|^q$  is convex on  $[\kappa_1, \kappa_2]$  for some fixed  $q > 1$ , then we have the following inequality for generalized fractional integrals

$$\begin{aligned} & \left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} \left( \int_0^1 \Lambda(\xi)^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \\ & \quad + \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} \left( \int_0^1 \Delta(\xi)^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}} \end{aligned}$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ .

*Proof.* By the Lemma 3.1, we have

$$\begin{aligned} & \left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} \int_0^1 \Lambda(\xi) |f'(\xi\kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))| d\xi \\ & \quad + \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} \int_0^1 \Delta(\xi) |f'(\xi\kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))| d\xi. \end{aligned}$$





Using the Hölder’s inequality and the convexity of  $|f'|^q$ , we obtain

$$\begin{aligned}
 & \left| \frac{1}{\Lambda(0) + \Delta(0)} \times \left[ {}_{\kappa_2^-} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) + {}_{\kappa_1^+} I_{\varphi} f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\
 & \leq \frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Lambda(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 |f'(\xi\kappa_2 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))|^q d\xi \right)^{\frac{1}{q}} \\
 & + \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Delta(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 |f'(\xi\kappa_1 + (1 - \xi)(\kappa_1 + \kappa_2 - \kappa))|^q d\xi \right)^{\frac{1}{q}} \\
 & \leq \frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Lambda(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 [\xi |f'(\kappa_2)|^q + (1 - \xi) |f'(\kappa_1 + \kappa_2 - \kappa)|^q] d\xi \right)^{\frac{1}{q}} \\
 & + \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Delta(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 [\xi |f'(\kappa_1)|^q + (1 - \xi) |f'(\kappa_1 + \kappa_2 - \kappa)|^q] d\xi \right)^{\frac{1}{q}} \\
 & = \frac{\kappa - \kappa_1}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Lambda(\xi))^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \\
 & + \frac{\kappa_2 - \kappa}{\Lambda(0) + \Delta(0)} \left( \int_0^1 (\Delta(\xi))^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}}.
 \end{aligned}$$

The proof is completed. □

**Corollary 4.9.** *Under assumption of Theorem 4.8 with  $\varphi(\xi) = \xi$ , we have the following inequality*

$$\begin{aligned}
 \left| \frac{1}{\kappa_2 - \kappa_1} \int_{\kappa_1}^{\kappa_2} f(\xi) d\xi - f(\kappa_1 + \kappa_2 - \kappa) \right| & \leq \frac{1}{(\kappa_2 - \kappa_1)(p + 1)^{\frac{1}{p}}} \left\{ (\kappa - \kappa_1)^2 \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \right. \\
 & \left. + (\kappa_2 - \kappa)^2 \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}} \right\}.
 \end{aligned}$$



**Remark 4.10.** Under assumption of Corollary 4.9 with  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$ , we have the following inequality

$$\begin{aligned} \left| \frac{1}{\kappa_2 - \kappa_1} \int_{\kappa_1}^{\kappa_2} f(\xi) d\xi - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| &\leq \frac{\kappa_2 - \kappa_1}{4(p+1)^{\frac{1}{p}}} \left[ \left( \frac{|f'(\kappa_1)|^q + 3|f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} + \left( \frac{|f'(\kappa_2)|^q + 3|f'(\kappa_1)|^q}{4} \right)^{\frac{1}{q}} \right] \\ &\leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{4}{p+1} \right)^{\frac{1}{p}} [|f'(\kappa_1)| + |f'(\kappa_2)|]. \end{aligned}$$

**Corollary 4.11.** Under assumption of Theorem 4.8 with  $\varphi(\xi) = \frac{\xi^\alpha}{\Gamma(\alpha)}$ , we have the following inequality

$$\begin{aligned} &\left| \frac{\Gamma(\alpha+1)}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \times \left[ J_{\kappa_2^-}^\alpha f(\kappa_1 + \kappa_2 - \kappa) + J_{\kappa_1^+}^\alpha f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ &\leq \frac{1}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \left( \frac{\alpha p}{\alpha p + 1} \right)^{\frac{1}{p}} \times \left\{ (\kappa - \kappa_1)^{\alpha+1} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \right. \\ &\quad \left. + (\kappa_2 - \kappa)^{\alpha+1} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 4.12.** Under assumption of Corollary 4.11 with  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$ , we have the following inequality

$$\begin{aligned} &\left| \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\kappa_2^-}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) + J_{\kappa_1^+}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \\ &\leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{\alpha p}{\alpha p + 1} \right)^{\frac{1}{p}} \times \left[ \left( \frac{|f'(\kappa_1)|^q + 3|f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} + \left( \frac{|f'(\kappa_2)|^q + 3|f'(\kappa_1)|^q}{4} \right)^{\frac{1}{q}} \right] \\ &\leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{4\alpha p}{\alpha p + 1} \right)^{\frac{1}{p}} [|f'(\kappa_1)| + |f'(\kappa_2)|]. \end{aligned}$$

**Corollary 4.13.** Under assumption of Theorem 4.8 with  $\varphi(\xi) = \frac{\xi^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ , we have the following inequality

$$\begin{aligned} &\left| \frac{\Gamma_k(\alpha+k)}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \times \left[ J_{\kappa_2^-,k}^\alpha f(\kappa_1 + \kappa_2 - \kappa) + J_{\kappa_1^+,k}^\alpha f(\kappa_1 + \kappa_2 - \kappa) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ &\leq \frac{1}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \left( \frac{\alpha p}{\alpha p + k} \right)^{\frac{1}{p}} \times \left\{ (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} \left[ \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right]^{\frac{1}{q}} \right. \\ &\quad \left. + (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} \left[ \frac{|f'(\kappa_1)|^q + |f'(\kappa_1 + \kappa_2 - \kappa)|^q}{2} \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 4.14.** Under assumption of Corollary 4.13 with  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$ , we have the following inequality

$$\left| \frac{2^{\frac{\alpha}{k}-1} \Gamma_k(\alpha+k)}{(\kappa_2 - \kappa_1)^{\frac{\alpha}{k}}} \left[ J_{\kappa_2^-,k}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) + J_{\kappa_1^+,k}^\alpha f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{(\kappa_2 - \kappa_1)}{4} \left( \frac{4\alpha p}{\alpha p + k} \right)^{\frac{1}{p}} [|f'(\kappa_1)| + |f'(\kappa_2)|].$$



**Theorem 4.15.**  $f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(\kappa_1, \kappa_2)$  with  $0 \leq \kappa_1 < \kappa_2$ . If  $|f'|$  is convex on  $[\kappa_1, \kappa_2]$ , then we have the following inequality for generalized fractional integrals

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \times \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{\Psi(1) + \Theta(1)} \times \left\{ (\kappa - \kappa_1) \left| f'(\kappa_2) \right| \int_0^1 (1 - \xi) \Psi(\xi) d\xi + (\kappa_2 - \kappa) \left| f'(\kappa_1) \right| \int_0^1 (1 - \xi) \Theta(\xi) d\xi \right. \\ & \quad \left. + \left| f'(\kappa_1 + \kappa_2 - \kappa) \right| \int_0^1 \xi [(\kappa - \kappa_1) \Psi(\xi) + (\kappa_2 - \kappa) \Theta(\xi)] d\xi \right\} \end{aligned} \tag{4.5}$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\Psi$  and  $\Theta$  are defined as in (3.4).

*Proof.* By the Lemma 3.2, we have

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \times \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{(\kappa - \kappa_1)}{\Psi(1) + \Theta(1)} \left| \int_0^1 \Psi(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2) d\xi \right| \\ & \quad + \frac{(\kappa_2 - \kappa)}{\Psi(1) + \Theta(1)} \left| \int_0^1 \Theta(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1) d\xi \right| \\ & \leq \frac{(\kappa - \kappa_1)}{\Psi(1) + \Theta(1)} \int_0^1 |\Psi(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2)| d\xi \\ & \quad + \frac{(\kappa_2 - \kappa)}{\Psi(1) + \Theta(1)} \int_0^1 |\Theta(\xi) f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1)| d\xi. \end{aligned}$$

Using the convexity of  $|f'|$ , we obtain

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \times \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{(\kappa - \kappa_1)}{\Psi(1) + \Theta(1)} \left\{ \int_0^1 \Psi(\xi) [\xi |f'(\kappa_1 + \kappa_2 - \kappa)| + (1 - \xi) |f'(\kappa_2)|] d\xi \right. \\ & \quad \left. + \frac{(\kappa_2 - \kappa)}{\Psi(1) + \Theta(1)} \int_0^1 \Theta(\xi) [\xi |f'(\kappa_1 + \kappa_2 - \kappa)| + (1 - \xi) |f'(\kappa_1)|] d\xi \right\} \\ & = \frac{1}{\Psi(1) + \Theta(1)} \times \left\{ (\kappa - \kappa_1) |f'(\kappa_2)| \int_0^1 (1 - \xi) \Psi(\xi) d\xi + (\kappa_2 - \kappa) |f'(\kappa_1)| \int_0^1 (1 - \xi) \Theta(\xi) d\xi \right. \\ & \quad \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \int_0^1 \xi [(\kappa - \kappa_1) \Psi(\xi) + (\kappa_2 - \kappa) \Theta(\xi)] d\xi \right\}. \end{aligned}$$

This completes the proof. □

**Remark 4.16.** If we choose  $\varphi(\xi) = \xi$  in Theorem 4.15, then the inequality (4.5) reduces to the inequality (4.3).



**Corollary 4.17.** *Under assumption of Theorem 4.15 with  $\varphi(\xi) = \frac{\xi^\alpha}{\Gamma(\alpha)}$ , we have the following inequality*

$$\begin{aligned} & \left| \frac{\Gamma(\alpha + 1)}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \times \left[ J_{(\kappa_1 + \kappa_2 - \kappa)^-}^\alpha f(\kappa_2) + J_{(\kappa_1 + \kappa_2 - \kappa)^+}^\alpha f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \frac{1}{\alpha + 2} \times \left\{ \frac{1}{(\alpha + 1)} \left[ (\kappa - \kappa_1)^{\alpha+1} |f'(\kappa_2)| + (\kappa_2 - \kappa)^{\alpha+1} |f'(\kappa_1)| \right] \right. \\ & \quad \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \left[ (\kappa_2 - \kappa)^{\alpha+1} + (\kappa - \kappa_1)^{\alpha+1} \right] \right\}. \end{aligned}$$

**Remark 4.18.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.17, then we have the following inequality

$$\left| \frac{2^{\alpha-1} \Gamma(\alpha + 1)}{(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^-}^\alpha f(\kappa_2) + J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^+}^\alpha f(\kappa_1) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{(\kappa_2 - \kappa_1)}{4(\alpha + 1)} [|f'(\kappa_1)| + |f'(\kappa_2)|]$$

which is given by Sarikaya and Yildirim in [36].

**Corollary 4.19.** *Under assumption of Theorem 4.15 with  $\varphi(\xi) = \frac{\xi^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ , we have the following inequality*

$$\begin{aligned} & \left| \frac{\Gamma_k(\alpha + k)}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \times \left[ J_{(\kappa_1 + \kappa_2 - \kappa)^-, k}^\alpha f(\kappa_2) + J_{(\kappa_1 + \kappa_2 - \kappa)^+, k}^\alpha f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^{\alpha/k} + (\kappa - \kappa_1)^{\alpha/k}} \frac{k}{(\alpha + 2k)} \\ & \quad \times \left\{ \frac{k}{(\alpha + k)} \left[ (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} |f'(\kappa_2)| + (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} |f'(\kappa_1)| \right] \right. \\ & \quad \left. + |f'(\kappa_1 + \kappa_2 - \kappa)| \left[ (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} + (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} \right] \right\}. \end{aligned}$$

**Remark 4.20.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.19, then we have the following inequality

$$\left| \frac{2^{\frac{\alpha}{k}-1} \Gamma_k(\alpha + k)}{(\kappa_2 - \kappa_1)^{\frac{\alpha}{k}}} \left[ J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^-, k}^\alpha f(\kappa_2) + J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^+, k}^\alpha f(\kappa_1) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \leq \frac{(\kappa_2 - \kappa_1)k}{4(\alpha + k)} [|f'(\kappa_1)| + |f'(\kappa_2)|]$$

which is given by Farid et al. in [15].

**Theorem 4.21.**  *$f : [\kappa_1, \kappa_2] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(\kappa_1, \kappa_2)$  with  $0 \leq \kappa_1 < \kappa_2$ . If  $|f'|^q$  is convex on  $[\kappa_1, \kappa_2]$  for some fixed  $q > 1$ , then we have the following inequality for generalized fractional integrals*

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \tag{4.6} \\ & \leq \frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} \left( \int_0^1 \Psi(\xi)^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \\ & \quad + \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} \left( \int_0^1 \Theta(\xi)^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}} \end{aligned}$$

with all  $\kappa \in [\kappa_1, \kappa_2]$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ .



*Proof.* By the Lemma 3.2, we have

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} \int_0^1 \Psi(\xi) |f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2)| d\xi \\ & \quad + \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} \int_0^1 \Theta(\xi) |f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1)| d\xi. \end{aligned}$$

Using the Hölder’s inequality and the convexity of  $|f'|^q$ , we obtain

$$\begin{aligned} & \left| \frac{1}{\Psi(1) + \Theta(1)} \left[ {}_{(\kappa_1 + \kappa_2 - \kappa)^-} I_\varphi f(\kappa_2) + {}_{(\kappa_1 + \kappa_2 - \kappa)^+} I_\varphi f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Psi(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 |f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_2)|^q d\xi \right)^{\frac{1}{q}} \\ & \quad + \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Theta(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 |f'(\xi(\kappa_1 + \kappa_2 - \kappa) + (1 - \xi)\kappa_1)|^q d\xi \right)^{\frac{1}{q}} \\ & \leq \frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Psi(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 [(1 - \xi)|f'(\kappa_2)|^q + \xi|f'(\kappa_1 + \kappa_2 - \kappa)|^q] d\xi \right)^{\frac{1}{q}} \\ & \quad + \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Theta(\xi))^p d\xi \right)^{\frac{1}{p}} \times \left( \int_0^1 [(1 - \xi)|f'(\kappa_1)|^q + \xi|f'(\kappa_1 + \kappa_2 - \kappa)|^q] d\xi \right)^{\frac{1}{q}} \\ & = \frac{\kappa - \kappa_1}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Psi(\xi))^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \\ & \quad + \frac{\kappa_2 - \kappa}{\Psi(1) + \Theta(1)} \left( \int_0^1 (\Theta(\xi))^p d\xi \right)^{\frac{1}{p}} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}}. \end{aligned}$$

The proof is completed. □

**Remark 4.22.** If we choose  $\varphi(\xi) = \xi$  in Theorem 4.21, then inequality (4.7) reduces to the inequality (4.5).



**Corollary 4.23.** Under assumption of Theorem 4.21 with  $\varphi(\xi) = \frac{\xi^\alpha}{\Gamma(\alpha)}$ , we have the following inequality

$$\begin{aligned} & \left| \frac{\Gamma(\alpha+1)}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \times [J_{(\kappa_1 + \kappa_2 - \kappa)^-}^\alpha f(\kappa_2) + J_{(\kappa_1 + \kappa_2 - \kappa)^+}^\alpha f(\kappa_1)] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^\alpha + (\kappa - \kappa_1)^\alpha} \left( \frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \times \left\{ (\kappa - \kappa_1)^{\alpha+1} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + (\kappa_2 - \kappa)^{\alpha+1} \left( \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_1)|^q}{2} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 4.24.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.23, then we have the following inequality

$$\begin{aligned} & \left| \frac{2^{\alpha-1} \Gamma(\alpha+1)}{(\kappa_2 - \kappa_1)^\alpha} \left[ J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^-}^\alpha f(\kappa_2) + J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^+}^\alpha f(\kappa_1) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \tag{4.7} \\ & \leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \times \left\{ \left( \frac{|f'(\kappa_1)|^q + 3|f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} + \left( \frac{3|f'(\kappa_1)|^q + |f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} \right\} \\ & \leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{4}{\alpha p + 1} \right)^{\frac{1}{p}} [|f'(\kappa_1)| + |f'(\kappa_2)|] \end{aligned}$$

which is proved by Sarikaya and Yildirim in [36].

**Corollary 4.25.** Under assumption of Theorem 4.21 with  $\varphi(\xi) = \frac{\xi^{\frac{\alpha}{k}}}{k\Gamma_k(\alpha)}$ , we have the following inequality

$$\begin{aligned} & \left| \frac{\Gamma_k(\alpha+k)}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \times \left[ J_{(\kappa_1 + \kappa_2 - \kappa)^-,k}^\alpha f(\kappa_2) + J_{(\kappa_1 + \kappa_2 - \kappa)^+,k}^\alpha f(\kappa_1) \right] - f(\kappa_1 + \kappa_2 - \kappa) \right| \\ & \leq \frac{1}{(\kappa_2 - \kappa)^{\frac{\alpha}{k}} + (\kappa - \kappa_1)^{\frac{\alpha}{k}}} \left( \frac{k}{\alpha p + k} \right)^{\frac{1}{p}} \times \left\{ (\kappa - \kappa_1)^{\frac{\alpha}{k}+1} \left[ \frac{|f'(\kappa_1 + \kappa_2 - \kappa)|^q + |f'(\kappa_2)|^q}{2} \right]^{\frac{1}{q}} \right. \\ & \quad \left. + (\kappa_2 - \kappa)^{\frac{\alpha}{k}+1} \left[ \frac{|f'(\kappa_1)|^q + |f'(\kappa_1 + \kappa_2 - \kappa)|^q}{2} \right]^{\frac{1}{q}} \right\}. \end{aligned}$$

**Remark 4.26.** If we choose  $\kappa = \frac{\kappa_1 + \kappa_2}{2}$  in Corollary 4.25, then we have the following inequality

$$\begin{aligned} & \left| \frac{2^{\frac{\alpha}{k}-1} \Gamma_k(\alpha+k)}{(\kappa_2 - \kappa_1)^{\frac{\alpha}{k}}} \left[ J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^-,k}^\alpha f(\kappa_2) + J_{\left(\frac{\kappa_1 + \kappa_2}{2}\right)^+,k}^\alpha f(\kappa_1) \right] - f\left(\frac{\kappa_1 + \kappa_2}{2}\right) \right| \\ & \leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{k}{\alpha p + k} \right)^{\frac{1}{p}} \times \left\{ \left( \frac{|f'(\kappa_1)|^q + 3|f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} + \left( \frac{3|f'(\kappa_1)|^q + |f'(\kappa_2)|^q}{4} \right)^{\frac{1}{q}} \right\} \\ & \leq \frac{\kappa_2 - \kappa_1}{4} \left( \frac{4k}{\alpha p + k} \right)^{\frac{1}{p}} [|f'(\kappa_1)| + |f'(\kappa_2)|]. \end{aligned}$$

which is given by Farid et al. in [15].

## 5. CONCLUSIONS

In this study, by obtained fractional equalities, we establish some midpoint type inequalities via generalized fractional integrals for functions whose derivatives in absolute values are convex. In the future works, authors can prove similar inequalities by using the different kinds of convexities.



## REFERENCES

- [1] A. Abdeldaim, A. A. El-Deeb, P. Agarwal, and H. A. El-Sennary, *On some dynamic inequalities of Steffensen type on time scales*, Math. Methods Appl. Sci., *41*(12) (2018), 4737-4753.
- [2] P. Agarwal and J. E. Restrepo, *An extension by means of  $\omega$ -weighted classes of the generalized Riemann-Liouville  $k$ -fractional integral inequalities*, J. Math. Ineq., *14*(1) (2020), 35-46.
- [3] P. Agarwal, J. Tariboon, and S. K. Ntouyas, *Some generalized Riemann-Liouville  $k$ -fractional integral inequalities*, J. Inequal. Appl, *2016* (2016), 122.
- [4] M. A. Ali, H. Budak, and I. B. Sial, *Generalized fractional integral inequalities for product of two convex functions*, Italian J. Pure Appl. Math., *45* (2021), 689698.
- [5] M. A. Ali, H. Budak, M. Abbas, M. Z. Sarikaya, and A. Kashuri, *Hermite-Hadamard type inequalities for  $h$ -convex functions via generalized fractional integrals*, J. Math. Extension, *14*(4), (2020), 187-234.
- [6] M. Alomari, M. Darus, and U. S. Kirmaci, *Refinements of Hadamard-type inequalities for quasi-convex functions with applications to trapezoidal formula and to special means*, Comput. Math. Appl., *59*(1) (2010), 225-232.
- [7] G. A. Anastassiou, *General fractional Hermite-Hadamard inequalities using  $m$ -convexity and  $(s, m)$ -convexity*, Frontiers in Time Scales and Inequalities, *237* (2016), 237-255.
- [8] A. G. Azpeitia, *Convex functions and the Hadamard inequality*, Rev. Colombiana Math., *28* (1994), 7-12.
- [9] H. Budak and P. Agarwal, *New generalized midpoint type inequalities for fractional integral*, Miskolc Math. Notes, *20*(2) (2019), 781-793.
- [10] H. Budak, F. Ertuğral, and E. Pehlivan, *Hermite-Hadamard type inequalities for twice differentiable functions via generalized fractional integrals*, Filomat, *33*(15) (2019), 4967-4979.
- [11] J. de la Cal, J. Carcamob, and L. Escauriaza, *A general multidimensional Hermite-Hadamard type inequality*, J. Math. Anal. Appl., *356* (2009), 659-663.
- [12] H. Chen and U.N. Katugampola, *Hermite-Hadamard and Hermite-Hadamard-Fejér type inequalities for generalized fractional integrals*, J. Math. Anal. Appl., *446*(2) (2017), 1274-1291.
- [13] S.S. Dragomir and C.E.M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000. Online:[<http://rgmia.org/papers/monographs/Master2.pdf>].
- [14] S.S. Dragomir and R.P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. lett., *11*(5) (1998), 91-95.
- [15] G. Farid, A. Ur Rehman, and M. Zahra, *On Hadamard type inequalities for  $k$ -fractional integrals*, Konurap J. Math., *4*(2) (2016), 79-86.
- [16] G. Farid; A. Ur Rehman and Q. Ul Ain,  *$K$ -fractional integral inequalities of Hadamard type for  $(h - m)$ -convex functions*, Comput. Methods Differ. Equ., *8*(1) (2020), 119-140.
- [17] M. Iqbal, M.I. Bhatti, and K. Nazeer, *Generalization of Inequalities Analogous to Hermite-Hadamard Inequality via Fractional Integrals*, Bull. Korean Math. Soc., *52*(3) (2015), 707716.
- [18] İ. İscan and S. Wu, *Hermite-Hadamard type inequalities for harmonically convex functions via fractional integrals*, Appl. Math. Comput., *238*(1) (2014), 237-244.
- [19] İ. İscan, *On generalization of different type integral inequalities for  $s$ -convex functions via fractional integrals*, Math. Sci. Appl. E Notes, *2*(1), 2014, 55-67.
- [20] M. Jleli and B. Samet, *On Hermite-Hadamard type inequalities via fractional integrals of a function with respect to another function*, J. Nonlinear Sci. Appl., *9*(3) (2016), 1252-1260.
- [21] D. Kaur, P. Agarwal, M. Rakshit and M. Chand, *Fractional calculus involving  $(p, q)$ -mathieu type series*, Appl. Math. Nonlinear Sci., *5*(2) (2020), 15-34.
- [22] R. Khalil, M. Al Horani, A. Yousef, and M. Sababheh, *A new definition of fractional derivative*, J. Comput. Appl. Math., *264* (2020), 65-70.
- [23] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204, Elsevier Sci. B.V., Amsterdam, 2006.
- [24] U. S. Kirmaci, *Inequalities for differentiable mappings and applications to special means of real numbers to midpoint formula*, Appl. Math. Comput., *147*(1) (2004), 137-146.



- [25] X. Liu, L. Zhang, P. Agarwal, and G. Wang, *On some new integral inequalities of Gronwall–Bellman–Bihari type with delay for discontinuous functions and their applications*, *Indag. Math.*, *27*(1) (2016), 1-10.
- [26] K. Mehrez and P. Agarwal, *New Hermite–Hadamard type integral inequalities for convex functions and their applications*, *J. Comput. Appl. Math.*, *350* (2020), 274-285.
- [27] S. Mubeen and G. M Habibullah, *k-Fractional integrals and application*, *Int. J. Contemp. Math. Sciences*, *7*(2) (2012), 89-94.
- [28] M. A. Noor and M. U. Awan, *Some integral inequalities for two kinds of convexities via fractional integrals*, *TJMM*, *5*(2) (2013), 129-136.
- [29] J.E. Pečarić, F. Proschan, and Y.L. Tong, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, Boston, 1992.
- [30] M. E. Özdemir, M. Avci-Ardıç, and H. Kavurmacı-Önalın, *Hermite-Hadamard type inequalities for s-convex and s-concave functions via fractional integrals*, *Turkish J. Science*, *1*(1) (2016), 28-40.
- [31] M. E. Özdemir, M. Avci, and E. Set, *On some inequalities of Hermite–Hadamard-type via m-convexity*, *Appl. Math. Lett.*, *23*(9) (2010), 1065–1070.
- [32] M. E. Özdemir, M. Avci, and H. Kavurmaci, *Hermite–Hadamard-type inequalities via  $(\alpha, m)$ -convexity*, *Comput. Math. Appl.*, *61*(9) (2011), 2614–2620.
- [33] S. Qaisar and S. Hussain, *On Hermite-Hadamard type inequalities for functions whose first derivative absolute values are convex and concave*, *Fasc. Math.*, *58* (2017), 155–166.
- [34] A. Saglam, M. Z. Sarikaya, and H. Yildirim, *Some new inequalities of Hermite-Hadamard's type*, *Kyungpook Math. J.*, *50*(3) (2010), 399-410.
- [35] M. Z. Sarikaya and F. Ertuğral, *On the generalized Hermite-Hadamard inequalities*, *An. Univ. Craiova Ser. Mat. Inform.*, *47*(1) (2020), 193-213.
- [36] M. Z. Sarikaya and H. Yildirim, *On Hermite-Hadamard type inequalities for Riemann-Liouville fractional integrals*, *Miskolc Math. Notes*, *17*(2) (2016), 1049–1059.
- [37] M. Z. Sarikaya, E. Set, H. Yaldiz, and N., Basak, *Hermite -Hadamard's inequalities for fractional integrals and related fractional inequalities*, *Math. Comput. Model.*, *57*(9-10) (2013), 2403–2407.
- [38] M. Z. Sarikaya and H. Budak, *Generalized Hermite-Hadamard type integral inequalities for fractional integrals*, *Filomat*, *30*(5) (2016), 1315–1326.
- [39] M. Z. Sarikaya, A. Akkurt, H. Budak, M. E. Yildirim, and H. Yildirim, *Hermite-Hadamard's inequalities for conformable fractional integrals*, *Int. J. Optim. Control. Theor. Appl.*, *9*(1) (2019), 49-59.
- [40] E. Set, M. E. Ozdemir, and M. Z. Sarikaya, *New inequalities of Ostrowski's type for s-convex functions in the second sense with applications*, *Facta Universitatis, Ser. Math. Inform.*, *27*(1) (2012), 67-82.
- [41] E. Set, M. Z. Sarikaya, M. E. Ozdemir, and H. Yildirim, *The Hermite-Hadamard's inequality for some convex functions via fractional integrals and related results*, *J. Appl. Math. Stat. Inform.*, *10*(2) (2014), 69-83.
- [42] E. Set, A. Ocak Akdemir, and F. Ozata, *Grüss type inequalities for fractional integral operator involving the extended generalized Mittag-Leffler function*, *Appl. Comput. Math.*, *19*(3) (2020), 422-434.
- [43] J. Wang, X. Li, M. Fečkan, and Y. Zhou, *Hermite–Hadamard-type inequalities for Riemann–Liouville fractional integrals via two kinds of convexity*, *Appl. Anal.*, *92*(11) (2012), 2241–2253.
- [44] J. Wang, X. Li, and C. Zhu, *Refinements of Hermite-Hadamard type inequalities involving fractional integrals*, *Bull. Belg. Math. Soc. Simon Stevin*, *20*(4) (2013), 655–666.
- [45] M. F. Yassen, A. A. Attiya, and P. Agarwal, *Subordination and Superordination properties for certain family of analytic functions associated with Mittag-Leffler function*, *Symmetry*, *12*(10) (2020), 17-24.
- [46] Y. Zhang and J. Wang, *On some new Hermite-Hadamard inequalities involving RiemannLiouville fractional integrals*, *J. Inequal. Appl.*, *2013* (2013), 220.

