



## **Review Article**

### **Zoonotic diseases associated with pet birds**

**Seyedeh Alemeh Hosseinian\***

Avian Disease Research Center, Department of Clinical Science, School of Veterinary Medicine, Shiraz University, Shiraz, Iran

\***Corresponding author:** [s-hosseinian@shirazu.ac.ir](mailto:s-hosseinian@shirazu.ac.ir)

(Received 23 September 2022, Accepted 16 October 2022)

#### **Summary**

Birds are the most popular pet animals kept by humans in many areas of the world. Budgerigars, Canaries, Lovebirds, and Cockatiels are the most common pet birds. Zoonotic diseases are one of the most critical concerns related to pet birds worldwide. The people at the most risk of zoonoses are immunocompromised patients, veterinarians, pet bird owners, pet bird shops, and workers in pet exhibitions. Zoonotic diseases are transmitted via the fecal-oral, inhalation, and vector-borne routes. Zoonotic pathogens infect humans through direct or indirect contact with infected birds, contaminated birds' cages, food dishes, or droppings. Several bacterial, viral, parasitic, and fungal zoonoses are known in pet birds. Some zoonoses, such as highly pathogenic avian influenza, salmonellosis, and chlamydiosis, have significant public health risks that lead to serious human diseases. Newcastle disease and giardiasis are minor zoonoses that cause self-limiting infections in humans. Salmonellosis and campylobacteriosis are food-borne zoonoses with global importance in human health. Biosecurity and hygiene, cleaning and disinfecting bird cages, and using human protective equipment can help to control zoonoses in birds and humans.

**Keywords:** Pet birds, Prevention, Public health, Zoonotic pathogens, Zoonoses.

#### **Introduction**

Keeping animals and birds as pets have been trending over the past several decades. Birds are the most popular pets for their colorful and attractive appearance (Chomel and Sun, 2011). According to the statistics, birds are the third and fourth most popular companion animal to keep as a pet in the European Union (EU) and United States, respectively (Peng and Broom, 2021). A pet or companion bird is a bird kept and bred primarily for ornamental purposes (Boseret et al., 2013). The three most common orders of birds kept as pet birds

include Psittaciformes (parakeets, parrots, love birds, cockatiel, and budgerigars), Passeriformes (finch, canary, mynah, starling, and sparrows), and Columbiformes (pigeons and doves). Budgerigars (*Melopsittacus undulatus*), canaries (*Serinus canaria*), lovebirds (*Agapornis sp.*), and cockatiels (*Nymphicus hollandicus*) are the most common pet birds in developed and developing countries (Boseret et al., 2013; Peng and Broom, 2021). Zoonotic diseases are one of the most critical concerns related to pet birds (Rahman et al., 2020). Zoonoses are infectious diseases transmitted

between animals or birds and humans. Zoonotic diseases transmit directly or indirectly from birds to humans (Guo et al., 2022). Some zoonoses transmit via asymptomatic reservoirs such as wild birds (Rahman et al., 2020). Some zoonoses, such as highly pathogenic avian influenza, salmonellosis, and chlamydiosis, are significant public health concerns leading to serious human diseases (Boseret et al., 2013). Salmonellosis and campylobacteriosis are food-borne zoonoses with global importance on human health, which can be transmitted from pet birds to humans (Rahman et al., 2020).

The severity of zoonotic diseases depends on various factors, such as organism's virulence, route of transmission or infection, and human's immunity level. Children, older people, immunocompromised and cancer patients are more likely to get zoonotic diseases (Pfeffer and Dobler, 2010). The people at the most risk of zoonoses are veterinarians, breeders of ornamental birds, pet birds owners, pet bird shops, and workers in exhibitions that have close contact with pet birds (Boseret et al., 2013). This review briefly discusses the most important zoonotic diseases transmitted from pet birds, their transmission route, and guidelines to control and prevent transmission from birds to humans.

### **Zoonoses transmission route**

The infectious pathogens of zoonoses disseminate through various routes. The pathogens are excreted from infected birds' feathers, feces, and various secretions (like saliva, ocular, tracheal, and other body fluids). Infected feathers, feces, and secretions infect food dishes, water bowls, and the environment in which birds live (including cages and soil). Infectious pathogens may persist in infected feces and soil for a long time (Hollingsworth et al., 2015). The zoonotic agents can be transmitted directly (through direct contact with sick birds) or indirectly (through touch with contaminated cages, food dishes, water bottles, and the environment around birds) to humans (Rahman et al., 2020). Zoonotic pathogens can enter the human body via various methods, including

ingestion, inhalation, mucous membrane, and damaged skin (Heesterbeek et al., 2015).

Some birds, particularly wild birds, can be asymptomatic reservoirs for zoonotic organisms and transmit them to humans without having any clinical signs. Various pathogens may be transferred from wild birds to pet birds kept outdoors (via contaminated feces), and humans become infected following contact with these pet birds (Ebani et al., 2021). Mites, mosquitoes, and ticks are mechanical vectors for some zoonotic pathogens, and the infectious agents are transmitted to humans via biting (Socha et al., 2022). Bird fairs, live pet shops, and markets are potential sources for some important zoonotic diseases, such as chlamydiosis and avian influenza (Magouras et al., 2020). Some zoonoses are food-borne diseases and can be transmitted from food chains to humans. *Salmonella* spp. is shed from sick birds' excreta and can contaminate birds' eggs and meat, and contaminated eggs and meat infect humans (Ehuwa et al., 2021). In zoonotic diseases, mostly transmission happens from infected birds to humans, but transmission between humans (inter-human transfer) is usually rare (Heesterbeek et al., 2015).

### **Classification of zoonoses**

Zoonoses are classified depending on the etiology of infection into various groups: bacterial, viral, parasitic, and fungal zoonoses (Chomel, 2009). Table 1 summarizes the most important zoonotic disease with their etiological agents, clinical signs, and clinical symptoms reported in humans.

#### ***Bacterial zoonoses***

Various bacterial species cause bacterial zoonosis. The most common zoonotic bacteria transmitted from birds to humans include *Salmonella* spp., *Chlamydia* spp., *Mycobacterium* spp., *Listeria* spp., *Escherichia coli*, *Campylobacter* spp., and *Staphylococcus* spp. (Day et al., 2016; Boroomand and Faryabi, 2020). Some zoonotic bacterial diseases, such as chlamydiosis and salmonellosis, have a higher occurrence in pet birds, which may be transferred to their owners leading to infection in humans (Jacob and Lorber, 2015). The bacterial

zoonotic agents with low occurrence in pet birds are *Yersinia* spp., *Pasteurella* spp., *Mycoplasma* spp., and *Klebsiella* spp. (Dorrestein et al., 2009; Jorn et al., 2009). *Escherichia coli* O157:H7, *Salmonella* spp., and *Mycobacterium Avium* can transmit from passerine birds (such as starlings) to

humans (Gaukler et al., 2009; Kauman and LeJeune, 2011). Chlamydiosis, salmonellosis, mycobacteriosis, campylobacteriosis, and Lyme disease are the most critical bacterial zoonoses in pet birds (Rahman et al., 2020).

**Table 1.** Summary of the most important zoonotic diseases in pet birds (Boseret et al., 2013; Contreras et al., 2016).

| Disease            | Etiology   | Sensitive host                | Transmission route to humans  | Major clinical signs in birds   | Major Clinical symptoms in human  | Risk factors  | OIE listed disease |
|--------------------|--|-------------------------------|---|---|---|---|--------------------|
| Histoplasmosis     | <i>Histoplasma capsulatum</i>                                  | All                           | Airborne, Inhalation of the fungus  | Often asymptomatic, chronic gastrointestinal infection and disseminated infections  | Often asymptomatic, weight loss, respiratory symptoms chronic progressive lung disease, chronic cutaneous, systemic disease, hepatosplenomegaly, and hematologic disturbances   | Immunocompromised individuals, Caves, guano   | No                 |
| Dermatophytosis    | <i>Microsporum</i> spp.,<br><i>Trichophyton</i> spp.           | All                           | Direct contact with infected birds or material                                    | Classical ring lesion with central healing and crusts at the peripheral area, some degree of folliculitis   | Skin lesions as red areas, along with skin itching and central pallor   | Immunocompromised individuals   | No                 |
| Giardiasis         | <i>Giardia</i> spp.  | All                           | Fecal-oral  | Asymptomatic and diarrhea   | Sometimes asymptomatic weight loss, diarrhea, and abdominal pain  | Lack of hygiene in bird equipment, water and feeders  | No                 |
| Cryptosporidiosis  | <i>Cryptosporidium</i> spp.                                    | All                           | Fecal-oral  | Asymptomatic; diarrhea, respiratory symptoms  | Sometimes asymptomatic, slight fever, diarrhea, abdominal pain, nausea and malaise; organ lesions (liver, pancreas, and respiratory tract)  | Lack of hygiene in bird equipment, water and feeders  | No                 |
| Cryptococcosis     | <i>Cryptococcus neoformans</i> ,<br><i>Cryptococcus gattii</i> | Parrots, little pet birds     | Airborne, mainly by inhalation of fungus, occasionally through breaks in the skin | Asymptomatic, grows in the feces of pigeons and other bird species Focal or disseminated infection, affecting a single organ system or many, and central nervous system involvement | Mostly asymptomatic, fever, nausea, and vomiting respiratory and nervous symptoms, severe symptoms in immunocompromised individuals, Meningitis/meningoencephalitis<br>Cutaneous, ocular, pulmonary and central nervous system involvements | Aerosolized droppings   | No                 |
| Campylobacteriosis | <i>Campylobacter jejuni</i> ,<br><i>Campylobacter coli</i>     | Estrildidae mostly            | Food-borne  | Diarrhea in young birds, Inapparent in adults, and healthy carriers   | Acute gastrointestinal symptoms and Self-limiting infection   | High prevalence in poultry farms, Cross-contamination of chicken carcasses, and Lack of hygiene in food preparation | No                 |
| Tuberculosis       | <i>Mycobacterium avium</i> ,<br><i>M. tuberculosis</i>         | Psittacine birds and canaries | Fecal-oral, inhalation, and airborne  | Diarrhea, weight loss, granulomatous lesions in the gastrointestinal and respiratory tract  | Fever, abdominal pain, fatigue, chronic diarrhea, and anemia  | Respiratory secretions, Contaminated water, soil and dust, and Immunocompromised individuals                        | Yes                |

|                                      |  |   |   |  |   |  |     |
|--------------------------------------|--|---|---|--|---|--|-----|
| Salmonellosis                        | <i>Salmonella</i> spp.                 | All (open-air aviaries)                 | Foodborne, fecal-oral, contamination in the surroundings of companion birds | Diarrhea and mortality in young birds and inapparent in adult birds (Healthy carriers)                               | Fever, gastroenteritis, and abdominal pain  | Contamination of eggs, poultry meat and their by-products, lack of hygiene in food preparation, and little risk with companion birds | No  |
| Psittacosis/ chlamydiosis/ornithosis | <i>Chlamydia</i> spp.                  | Psittacine birds, canaries, and finches | Respiratory transmission (dried and aerosolized feces)                      | depression, anorexia, ruffled feathers, weight loss diarrhea, conjunctivitis, and serous ocular and nasal discharges | Cough, dyspnea, pleuritic chest pain, epistaxis, sore throat, hemoptysis, fever, malaise, anorexia, chills, nausea, vomiting, myalgias, arthralgias, headache, and abdominal pain | Companion birds (specially Psittacine birds) and Occupational zoonosis   | Yes |
| West Nile fever                      | flavivirus                             | All                                     | Percutaneous by mosquito bite ( <i>Culex</i> spp., <i>Aedes</i> spp.)       | Asymptomatic Occasionally, severe symptoms with high mortality   | Several neurological disorders, weakness, swollen lymph nodes, convulsions, tremors, disorientation, and neck stiffness, convulsions, and paralysis                               | Mosquito breeding sites and Sick or carrier wild birds   | Yes |
| Influenza disease                    | Orthomyxovirus                         | Passerines                              | Airborne, Secretions and feces of sick birds or healthy carriers            | High mortality in birds and Asymptomatic in wild carriers  | Respiratory symptoms High fatality rate   | Wild reservoir (waterfowl), Intermingling of wild birds of different origins, and Live bird markets                                  | Yes |
| Newcastle disease                    | Paramyxovirus                          | All                                     | Direct or indirect contact, respiratory secretions or droppings             | Usually respiratory symptoms, mild, moderate and very virulent forms   | Mild and self-limited conjunctivitis  | Direct contact with sick or carrier birds Live vaccines  | Yes |
| Lyme disease (borreliosis)           | <i>Borrelia burgdorferi</i> sensu lato | All                                     | Percutaneous (tick bite)  | Mainly asymptomatic  | Fever, headache, skin rash, and erythema migrans, and cardiac symptoms  | Migratory birds with <i>Ixodes</i> : geographical spread of <i>Borrelia burgdorferi</i>  | No  |

### Chlamydiosis

Chlamydiosis is one of the most important bacterial zoonoses caused by *Chlamydia psittaci* (Sachse et al., 2015). This organism is a small intracellular bacterium excreted from infected birds' nasal discharges and feces (Smith et al., 2011). Chlamydiosis is also known as psittacosis, ornithosis, or parrot fever (Cheong et al., 2019). The disease is called psittacosis if humans become infected via contact with psittacine birds. The term ornithosis is applied if the infection in humans happens due to contact with non-psittacine birds (Malik et al., 2021). All avian species are susceptible to chlamydiosis, but turkeys, pigeons, and ducks are the most vulnerable hosts for the disease. Pigeons, doves, waterfowl, herons, cockatoos, and parrots are the most commonly

represented sources of human infection (Circella et al., 2011; Sachse et al., 2015).

The infection is mainly transmitted by respiratory secretions and feces of sick birds. There is no report about the vertical transmission of *C. psittaci* through birds' eggs (Smith et al., 2011). The infected birds are a commonly asymptomatic carrier for *C. psittaci* but can shed the bacterium in feces and nasal discharges. Sometimes, non-specific clinical signs such as depression, anorexia, lethargy, ruffled feathers, and weight loss may be seen in some birds. Infected birds often develop diarrhea, conjunctivitis, and serous ocular and nasal discharge (Cheong et al., 2019). It is complicated to diagnose the disease, especially in asymptomatic birds. Tissue smears, feces, conjunctival, choanal, and cloacal swab samples, and liver biopsy can be used to diagnose infection

in birds. Various diagnostic tests, including antibody detection, antigen detection, bacteriologic culture, and polymerase chain reaction (PCR), are recommended for evidence of chlamydiosis in birds (Stokes et al., 2021).

Human infection with *C. psittaci* occurs by inhalation of infectious aerosols from dried feces, direct contact with a sick bird and their respiratory secretions, and the fecal-oral route (Malik et al., 2021). Usually, the incubation period in humans is 5 - 14 days. Flu-like symptoms such as fever, headache, dry cough, respiratory distress, muscle pain, and pneumonia are observed in infected people (Hogerwerf et al., 2017). Less common clinical manifestations in patients are hepatitis, arthritis, and meningoencephalitis. The infection severity in human varies from mild to sepsis with multiple organ failures (Smith et al., 2011). Disease in humans can be diagnosed using complement fixation test (CFT), enzyme-linked immunosorbent assay (ELISA), nucleic acid amplification tests (NAATs), and PCR (Bryan et al., 2019). Tetracycline, doxycycline, or azithromycin are drug choices. Treatment duration varies based on the drug type, but at least 14 days is recommended for tetracycline (Shi et al., 2021). Prevention of *C. psittaci* infection in humans is vital. Removing debris and fecal material from cages, and cleaning and disinfecting bird cages, water bowls, and food containers help avoid spreading infected material to humans. People should not work with large numbers of pet birds in closely and dusty areas. Also, people should use personal protective equipment that covers the head, eyes, and hands to prevent and control this disease (Malik et al., 2021).

### **Salmonellosis**

Salmonellosis is one of the most dangerous bacterial zoonoses with a global distribution caused by salmonella spp. (Barbour et al., 2015). It is one of the most prevalent food-borne diseases worldwide (Pui et al., 2011). Salmonellosis has remarkable diversity in the host range and is seen in humans, animals, and domestic and wild birds (Anamaria et al., 2018). About 2,500 Salmonella serovars have been identified worldwide, and the

best-known Salmonella serovars are *S. enterica*, which causes illness in birds, animals, and humans (El-Tayeb et al., 2017). *S. Pullorum* and *S. Gallinarum* are two of the most important serovars in poultry with considerable economic losses. *S. Pullorum* and *S. Gallinarum* are host-specific for birds and responsible for causing a wide range of infections (from subclinical to severe) in chickens (Andino and Hanning, 2015; Jajere, 2019). These serovars are non-pathogenic for humans and animals and have low zoonotic risk (Ren et al., 2017). *S. Typhimurium* and *S. Enteritidis* serovars are not host-specific and have a wide range of hosts. They are transmitted through food to humans and cause food poisoning and mass infection (Lawson et al., 2014).

Salmonellosis occurs in passerines, psittacines, turkeys, waterfowls, quails, pigeons, pheasants, parrots, and sparrows. The most susceptible pet birds to salmonella infection are pigeons, finches, canaries, and parrots. *S. Typhimurium* is frequently isolated from passerines, pigeons, ducks, and geese (Rahmani et al., 2011; Lawson et al., 2014). Salmonellosis is mainly an asymptomatic infection in pet birds. Some pet birds remain potential carriers for a long time and shed the bacteria into the environment through their feces (Wibisono et al., 2020). However, infected birds sometimes exhibit mild clinical signs, including depression, dehydration, anorexia, diarrhea, septicemia, and crop stasis. Granulomatous lesions in the spleen, ceca, and liver, ocular infection, arthritis, and osteomyelitis have been seen in infected passerines by *S. Typhimurium* (Boseret et al., 2013). The infection can be transmitted between birds horizontally (via direct and indirect contact with sick birds) and vertically (through eggs; Gow et al., 2009; Jones et al., 2011).

Salmonella transmission from birds to humans occurs via direct or indirect exposure to infected birds. Also, consuming water or food contaminated with infected birds' feces plays a significant role in human infection by Salmonella (Saravanan et al., 2015; Nidaullah et al., 2017). It has been reported that some outbreaks of *S. Typhimurium* in humans were following contact with the pellet food of

infected birds (Boseret et al., 2013). Furthermore, wild songbirds and starlings have been documented as potential *Salmonella* bacteria carriers to humans and mammals (Wibisono et al., 2020).

The prevalence of salmonellosis in humans has been increased in the 20<sup>th</sup> century due to the high consumption of animal products and poultry meat and eggs (Kallapura et al., 2014; Nidaullah et al., 2017). In addition, keeping pet birds and their close contact with humans is another reason for increasing the incidence of salmonellosis in humans in recent years (Antunes et al., 2016; Antillón et al., 2017). Eight to 48 hours after the *Salmonella* bacterium enters the human body, the disease appears in the form of gastroenterocolitis with clinical symptoms of increased body temperature, chill, vomiting, diarrhea, and myalgia in the infected individual. Vomiting, foul-smelling watery diarrhea, greenish color diarrhea, headache, abdominal cramps, fever, and nausea are more common clinical symptoms in infected humans (Jarvis et al., 2016; Antillón et al., 2017). Occasionally, human salmonellosis may be complicated by endocarditis, myelitis, and cerebral meningitis, requiring hospitalization. Usually, the illness in humans last 4 to 7 days, and death may rarely occur (Wu et al., 2011). Various reports indicate that in 2010 and 2011, about 9,000 cases of food poisoning caused by *Salmonella* were recorded in Poland, of which about 70% required hospitalization. Also, in 2013, about 8300 patients were identified in Poland, of which 70% were hospitalized. According to the European Food Safety Authority (EFSA) report, 93648 incidences of food poisoning by *Salmonella* were recorded in 30 EU countries in 2012. The EFSA also recorded 82694 cases in the EU in 2013 (Wojtyniak et al., 2013; Osek and Wiczorek, 2014).

The successful growth of organisms on selective enriched culture media is essential for definitive diagnosis. Sensitive antibiotics and supportive care are used for the treatment of infected humans. Disinfecting the surface and cage, cleaning the bird's water and food dishes, and avoiding contact with infected bird help prevent *Salmonella*

infection in humans (Hung et al., 2017; Kurtz and Goggins, 2017).

### *Mycobacteriosis*

*Mycobacteriosis* or tuberculosis is a significant disease caused by the bacterium *Mycobacterium avium*, which occurs in captive exotic, pet, domestic, and wild birds worldwide (Tsiouris et al., 2021). *M. avium* has four pathogenic subtypes for birds, including *M. avium* Subsp. *avium*, *M. avium* Subsp. *silvaticum*, *M. avium* Subsp. *hominissuis* (Haridy et al., 2014). *M. avium* Subsp. *avium* (serotypes 1, 2, and 3) is primary agent for avian mycobacteriosis (Haridy et al., 2014; Fulton et al., 2018). *M. avium* and *M. genavense* are the most common isolated species from pet birds, especially parrots, parakeets, and budgerigars (Lennox, 2007). *M. avium* Subsp. *paratuberculosis* is a pathogen of mammals (Tsiouris et al., 2021).

Avian tuberculosis affects all species of birds. Chickens, turkeys, pheasants, ducks, geese, and pet birds are the more susceptible host for tuberculosis (Fulton et al., 2018). Usually, mycobacteriosis affects birds older than one year. Infected birds with *M. avium* usually show no clinical symptoms but may become emaciated and lethargic. Diarrhea, weight loss, and colorless wattle and comb are seen in infected birds. Some birds develop respiratory distress, dyspnea, and sudden death due to granulomatous lesions in the lung. Granulomatous ocular lesions and skin lesions have also been reported in affected birds. *Mycobacteriosis* develops granulomatous lesions in various organs (including intestinal tract, liver, spleen, and lung) in birds (Dhama et al., 2011; Fulton et al., 2018). Infected, untreated, and carrier birds are potential reservoirs of bacteria for humans that have a significant impact on public health. *M. avium* is transmitted directly and indirectly to humans, which causes a progressive disease in humans, especially in immunosuppressive patients. Infected people show weight loss, fever, abdominal pain, fatigue, chronic diarrhea, and anemia. In some patients, localized diseases occur with central nervous system infection, endocarditis, and cervical lymphadenitis (Fulton et al., 2018).

*M. tuberculosis* is the main species responsible for tuberculosis in humans. Several reports documented that *M. tuberculosis* was isolated from canary and parrots in New York and Switzerland. These infected birds developed lethargy, osteomyelitis, and granulomatous hepatitis. It was reported that *M. tuberculosis* could circulate between humans and birds, which has public health importance. There is a concern that pet birds can be a potential reservoir of this organism for humans and other birds (Hoop, 2002; Steinmetz et al., 2006).

Identifying infected birds is one of the primary ways to prevent mycobacteriosis in humans. Mycobacterial culture, serological test, and PCR apply for the definitive diagnosis of mycobacterium in birds. The Tuberculin test is a more common serologic test for determining the prevalence of infection in birds. The tuberculin test is used for domestic fowl and is less useful in pet birds. In waterfowl birds, a whole blood stained-antigen agglutination test is used to diagnose mycobacteriosis (Tsiouris et al., 2021).

Antibiotics such as rifampin, ethambutol, azithromycin, or clarithromycin are used to treat mycobacteriosis in pet birds. Destroying bacteria with disinfectants and avoiding humans from contact with sick birds are the best ways to prevent mycobacterium infection in humans. Mycobacterium persists in the environment for several months. Therefore, disinfecting the environment and cage with bleach and the phenolic compound helps destroy and inactivate the bacteria (Fulton et al., 2018).

### ***Campylobacteriosis***

Campylobacteriosis is another dangerous zoonotic disease with significant veterinary and public health concerns caused by *Campylobacter* spp. (Kaakoush et al., 2015). Campylobacteriosis is one of the most important causes of global diarrheal illness (Hsieh and Sulaiman, 2018). *Campylobacter* spp. is a gram-negative bacterium with three strains: *Campylobacter jejuni*, *Campylobacter lari*, and *Campylobacter coli* (Maësaar et al., 2020). Poultry, turkey, chicken,

geese, and duck are infected with *Campylobacter* spp., particularly *C. jejuni* and *C. coli* (Kürekci et al., 2021). It has been reported that *C. jejuni* is the most isolated strain from broiler flocks (Maësaar et al., 2020). Wild and pet birds (primary canaries and parrots) are other asymptomatic reservoirs for *C. jejuni* (Mughini-Gras et al., 2021). *C. jejuni* and *C. coli* are colonized in the intestinal tract of chicken and turkey without any clinical disease. Wild birds carry these bacteria asymptotically, and their feces are a potential source for the organism (Antilles et al., 2015; Johansson et al., 2018). *C. jejuni* is the most common species (over 80%) isolated from human with campylobacteriosis (Taheri et al., 2019).

Campylobacteriosis is a food-borne disease, and consuming contaminated or undercooked meat is the primary transmission route from birds to humans (Mohan et al., 2013). Also, the infection is transmitted to humans through close exposure with infected birds or their feces and inhalation of the infected droplets and particles (Taff and Townsend, 2017). In addition, humans can be infected following a contact with surfaces and equipment contaminated with bird droppings (French et al., 2009; Mulder et al., 2020). After 2-5 days, the clinical symptoms develop in the form of fever, vomiting, watery or sticky diarrhea, abdominal pain, headache, and muscle pain in infected humans (Awofisayo-Okuyelu et al., 2017). The clinical infection mainly occurs in immunosuppressive people. Usually, the complication is rare; however, septicemia, endocarditis, and arthritis are sporadically seen. Meningitis, colitis, acute cholecystitis, and Guillain-Barré syndrome can rarely occur in patients (Fernandez-Cruz et al., 2010; Backert et al., 2017, Brooks et al., 2017). There are some reports of mild diarrhea in children infected by *C. lari* (Kaakoush et al., 2015). According to EFSA statistics, campylobacteriosis was a more common zoonosis in humans in 2010, and contaminated poultry meat was the primary source of *campylobacter* spp. (Kirk et al., 2010; Scallan et al., 2011).

Another serious public health concern of *Campylobacter* is antibiotic resistance, particularly for tetracyclines and fluoroquinolones. Ciprofloxacin resistance has been reported in some areas, such as the U.S. (Iannino et al., 2019). Histopathological examination, serologic tests, and PCR are used to diagnose the infection. Fluid and electrolyte therapy apply to the treatment of patients. Antibiotic treatment decreases the shed of *Campylobacterium* (Scallan et al., 2011).

#### ***Lyme disease***

Lyme disease (Lyme borreliosis) is a prevalent tick-borne infectious disease caused by the bacterium *Borrelia burgdorferi* (Hao et al., 2011). *Borrelia* spp. is transmitted to humans through the bite of an infected tick from the genus Ixodid (Ogden et al., 2015). The most common hosts for *Borrelia* spp. are wild birds, songbirds, sparrows, and canaries (Scott et al., 2012; Dumas et al., 2022). Birds can not directly transmit *Borrelia* spp. to humans but can carry ticks containing the organism to a new area. Therefore, migrating and wild birds have a significant role in the worldwide distribution of *Borrelia* spp. (Cohen et al., 2015; Dumas et al., 2022). The ticks are a vector for the organism transmitted from birds to humans by biting (Hamer et al., 2012).

Different species of *Borrelia*, including *B. burgdorferi*, *B. afzelii*, and *B. garinii*, were isolated from ticks collected on pet birds (Sürth et al., 2021). *B. garinii* is the most common isolated species of birds and the most common agent of Lyme disease in humans (Coipan et al., 2016; Barstad et al., 2018). Infected ticks attack the head and the skin around the ear and eyes of birds. Infected birds usually show no significant clinical signs but are a vector for infected ticks. Infected ticks may drop off the bird into a yard and garden, then they can later bite and infect a human. *Borrelia* spp. is entered into the body of humans through a bite from infected ticks. Fever, fatigue, headache, skin rash, and erythema migrans were seen in diseased humans (Stanek et al., 2012; Steere et al., 2016). However, the infection may spread to people's nervous systems, hearts, and joints. Diagnosis is based on clinical symptoms,

skin rash, and exposure to infected ticks (Mygland et al., 2009). Treatment is done with a few weeks of antibiotic therapy. Applying pesticides, removing ticks promptly, using insect repellent, and reducing tick habitat help prevent Lyme borreliosis (Steere et al., 2016).

#### ***Viral zoonoses***

Some viruses like avian influenza, avian paramyxovirus, and West Nile virus (WNV) are main viral zoonoses found in pet birds. Mostly, viral pathogens are transmitted between humans and birds through arthropod, mechanical and biological vector. Some zoonotic viruses are transmitted through wild birds as reservoirs (Venkatesan et al., 2010). Among zoonotic agents, avian influenza is the most dangerous pathogen. Infection with avian paramyxovirus in humans results in a self-limiting eye infection (Kozdruń et al., 2015).

#### ***Avian influenza***

The influenza virus belongs to the family Orthomyxoviridae. It has three types: A, B, and C. Influenza A virus is a zoonotic agent and infects a wide variety of hosts, including humans, sea mammals, pigs, horses, dogs, wild and domestic birds. Influenza B and C viruses naturally infect humans, and the influenza C virus has also been isolated from swine (Wright et al., 2013). Influenza A viruses are classified into various subtypes based on two surface glycoproteins: hemagglutinin (H) and neuraminidase (N) (Hill et al., 2019; Modiriamedan et al., 2021). Some subtypes are adapted only for birds and are not pathogenic for humans, known as the avian influenza virus (AIV) (Giotis et al., 2019). However, some other subtypes are not host-specific and infect various host ranges like birds, animals, and humans (Wang et al., 2021). Notably, all influenza A viruses have been isolated from wild birds, and birds are vital hosts for the ecology of the viruses (Zanaty et al., 2019). Infection with avian Influenza viruses (AIVs) in domestic poultry (turkey and chicken) can lead to respiratory diseases with various clinical symptoms from mild to severe (Karimi Madab et



al., 2013). The severity of clinical signs depends on the strain type of virus and host sensitivity (Samy et al., 2018). Many wild birds (including aquatic birds, free-ranging ducks, shorebirds, and passerines) are infected with all strains of influenza A (Bi et al., 2016). However, they are asymptomatic reservoirs for influenza A viruses and only shed viruses to the environment for a long time through their feces (Lang et al., 2016). Wild aquatic birds, especially waterfowl, can naturally transmit the infection to humans, domestic and pet birds (Ramey et al., 2022). Several types of psittacine birds, including lovebirds, parakeets, budgerigars, cockatiels, and parrots, are susceptible to AIV infection (Jiao et al., 2012a,b). Sometimes, the infected pet birds with AIV show no or few clinical signs (Su et al., 2015). There is a report of a 3-month-old red-lored Amazon parrot (*Amazona autumnalis*) infected with AIV (subtype H5N2) with clinical signs such as severe lethargy, melena, and regurgitation (Hawkins et al., 2006). The AIV transmits mainly between birds through fecal-oral and inhalation of infected droplets and aerosols route (Su et al., 2015). Some strains of influenza A (low pathogenic avian influenza virus) cause mild to moderate respiratory signs in broiler chickens. Other strains, termed highly pathogenic avian influenza, cause severe systemic infection with high mortality of about 100% in flock broilers (França and Brown, 2014.). Depression, anorexia, diarrhea, and neurologic signs are also seen in infected broilers with the Influenza virus. The principal diagnosis method is viral isolation from cloacal and tracheal swabs of sick birds (Spackman et al., 2014).

Human infection with some subtypes of AIV like H5N1, H7N9, and H9N2 has been reported (Lai et al., 2016). Transmission of AIV to humans occurs via direct or indirect contact with infected birds, oral-fecal, and inhalation of infected aerosols. It is noticeable that human-to-human transmission of avian influenza viruses is rare (Sergeev et al., 2013). Infected humans may show mild to severe clinical symptoms (Sun et al., 2010). Conjunctivitis, fever, cough, severe pneumonia, dyspnea, sepsis shock, and even death can develop

in infected people. Encephalitis and gastrointestinal symptoms have been reported in some people (Chen et al., 2014; Belser et al., 2018). It is noticeable that waterfowl strains of AIV have a low ability to replicate well in the human body and mainly immunosuppressive people are at risk for infection (Gray et al., 2011). The genome of the influenza virus is segmented; therefore, exchange of RNA segments between various subtypes can lead to producing a new virus that overcomes this host restriction (Long et al., 2019). It has been documented that human and avian influenza viruses can be well replicated in the pig's body. Pig is a mixing vessel for the genetic re-assortment between human and avian influenza viruses and the source of new viruses (Mostafa et al., 2018). Countries with close contact between humans, pigs, and birds can be considered an epicenter for emerging strains of influenza A virus. Therefore, pigs may transmit new viruses to birds (Zhang et al., 2021). Pet birds can be infected with new strains of influenza viruses via exposure to the feces of wild birds (Ebani et al., 2021). Live-bird markets and pet stores are important epicenters for all subtypes of influenza A and have an essential role in the transmission of virus to humans (Li et al., 2022).

Controlling the disease in poultry and other birds is the primary method of decreasing the risk to humans. Vaccinating chickens, avoiding human exposure to infected birds or their feces, and cleaning and disinfecting the cage and environment can help to prevent influenza infection in humans (Pan et al., 2022).

#### **West Nile Fever**

West Nile fever [known as West Nile fever virus (WNV)] is a viral disease in birds caused by the flavivirus genus (Chancey et al., 2015). The flavivirus is transmitted by arthropod vectors (mainly mosquitoes) (Michel et al., 2018). Canaries, parrots, and house sparrows are the primary hosts for WNV (Perez-Ramirez et al., 2014). Mostly, birds are infected subclinical and show no significant clinical signs. Sometimes, infected birds show ocular disease, convulsions, ataxia, torticollis, and paralysis of the pelvic and

thoracic limbs (David and Abraham, 2016). The WNV infection in some pet birds, such as psittacine and passerine birds, is fatal. Some pet birds survive and become potential reservoirs for human infections (Marzal et al., 2022). Wild, free-living, and migratory birds are also potential reservoirs for the virus (Ziegler et al., 2019). When the mosquito feeds on the sick bird, it becomes infected and then spreads the flavivirus to humans through biting (Kampen et al., 2020). The infection in birds is diagnosed by virus isolation and PCR. Infected tissues (spleen, brain, and kidneys) and cloacal or oral swabs of birds are used for virus isolation (Reemtsma et al., 2022).

Humans can be infected by flavivirus via direct or indirect exposure to infected birds, organ transplants, and blood donation products (Reemtsma et al., 2022). The virus causes respiratory signs in humans (Bai et al., 2019). The clinical symptoms in infected humans develop 3-14 days after biting an infected mosquito. Clinical symptoms are seen only in 20% of the infected humans, including fever, nausea, headache, vomiting, and rash (David and Abraham, 2016). Clinical signs may persist for a few days to a week. Severe clinical symptoms such as weakness, convulsions, tremors, disorientation, and neck stiffness may develop rarely. In these cases, the duration of the disease may last for weeks, and the neurological signs may be permanent (Bai et al., 2019; Reemtsma et al., 2022).

West Nile Fever treatment is primarily supportive. No treatment is needed in mild cases. Severe cases of infection in humans may require hospitalization. The primary way of prevention is to limit exposure to mosquitoes. Spraying the area with insecticides and using body protection clothes against the mosquito can help control the infection (Bai et al., 2019).

### ***Newcastle disease***

Newcastle disease is a minor zoonosis caused by an avian paramyxovirus of the family paramyxoviridae (Behboudi and Hamidi Sofiani, 2021). There are different paramyxovirus serotypes, and each serotype is adapted to a

specific host. Avian paramyxovirus serotype 1 (APMV-1) has significant economic and public health concerns, causing Newcastle disease in birds (Linde et al., 2011). Newcastle disease is a critical disease in poultry, leading to respiratory disease accompanied by a high mortality rate, intestinal tract, and neurological signs. Newcastle disease has high economic loss in the poultry industry worldwide due to weight loss, decreased weight gain, and increased mortality rate. Newcastle disease is notifiable and requires prompt steps for eradication (Narayanan et al., 2010).

Many species of birds, including poultry, pigeon, parrot, passerine, domestic and wild birds, are susceptible to APMV-1. Mostly, waterfowl birds are asymptomatic reservoirs and shed the APMV-1 via their feces (Munir et al., 2012). The virus is shed from respiratory secretions and feces of infected birds and can contaminate the environment. Direct or indirect contact with infected birds, environment, and equipment leads to transmitting the Newcastle disease between birds. The virus enters the bird's body due to inhalation and ingesting contaminated particles (Yune and Abdela, 2017). Serological tests, viral isolation, and molecular tests are used for Newcastle disease diagnosis in birds (Alazawy and Al Ajeeli, 2020).

APMV-1 is transmitted from birds to humans, causing mild flu-like symptoms, conjunctivitis, and laryngitis. Veterinarians, people working with live Newcastle virus vaccines, and people contacting sick birds are at risk of infection by APMV-1. In humans, clinical signs occur 1-2 days after infection. Infected humans show conjunctival redness, watery eyes, subconjunctival ecchymoses, and eyelid swelling. The illness is self-limiting, and symptoms usually persist for a few days (Ul-Rahman et al., 2022).

Pigeon paramyxovirus-1 (PPMV-1) is another zoonotic virus of the genus Avulavirus, family paramyxoviridae. Pigeons and doves are the primary hosts for PPMV-1. Infection with PPMV-1 leads to green diarrhea, lethargy, twisting of the neck, head flicking, and dyspnea in pigeons. Close contact with infected pigeons or their droppings

can transmit PPMV-1 to humans. The PPMV-1 infection causes respiratory disease in humans, especially in immunocompromised individuals (Kuiken et al., 2018)

### **Parasitic zoonoses**

There is a wide variety of cestodes, protozoa, trematodes, and nematodes, with zoonoses concern between animals and humans (Schurer et al., 2016). A few parasites like *Giardia* spp. and *Cryptosporidium* spp. in pet birds can lead to disease in humans. Parasites transfer to humans via various routes, including vectors, contact with infected birds, and exposure to contaminated birds' equipment and environments around the birds. Wild and pet birds can be potential reservoirs for the various parasites (Ebani et al., 2021).

#### ***Giardiasis***

Giardiasis is a zoonotic parasitic disease caused by *Giardia* spp. Different species of *Giardia* are known in birds: *G. psittaci*, *G. duodenalis*, and *G. ardeae* (Ryan and Cacciò, 2013). *G. psittaci* and *G. ardeae* are specific to birds and not transmissible to humans (Cano et al., 2016). *G. duodenalis* is a zoonotic parasite and usually causes a self-limiting infection in humans. It has been reported that parrots are the host for some species of *G. duodenalis*, which are colonized in the small intestine of birds and shed from their feces (da Cunha et al., 2017). Feces of various pet birds, mainly parrots, are a source of *Giardia* spp. infection in humans. Infection in humans is mainly asymptomatic, but sometimes has mild clinical signs like abdominal pain, diarrhea, and weight loss (Reboredo-Fernandez et al., 2015).

#### ***Cryptosporidiosis***

Cryptosporidiosis is caused by *Cryptosporidium* spp., which is isolated from droppings of various birds like pigeons, parrots, and songbirds. *C. galli*, *C. meleagridis*, and *C. baylei* are not zoonotic agents and only infect birds leading to respiratory, intestinal, and nephrotic clinical signs in the bird (Cano et al., 2016; da Cunha et al., 2017). *C. hominis* and *C. parvum* are two important *Cryptosporidium* spp. with zoonotic ability. These species are isolated from various types of pet birds,

which can lead to human infection (Braima et al., 2019). Contaminated areas, water, and food are sources of infection in humans. Very young children and HIV-positive patients are at the highest risk for cryptosporidiosis by *C. hominis* and *C. parvum*. Infected individuals develop respiratory signs and lesions in various organs such as gastrointestinal tract, liver, and pancreas (Sak et al., 2011; Braima et al., 2019).

#### ***Toxoplasmosis***

Toxoplasmosis is a zoonosis caused by the protozoan parasite *Toxoplasma gondii* infecting animals, birds, and mammals worldwide. *T. gondii* causes infection in humans, especially in immunocompromised individuals and unborn fetuses (Ammar et al., 2021). The most important hosts for the organism are finches, budgerigars, and canaries (Boseret et al., 2013). Huang et al. (2019) reported that 34.29% of Java sparrows (*Lonchura oryzivora*) are infected with *T. gondii*. In the previous studies, it has been reported that the prevalence of *T. gondii* in pigeons and pet parrots were 11.86% and 8.36%, respectively (Cong et al., 2012; Zhang et al., 2014). Exposure to infected feces or eating raw or undercooked meat has a vital role in the transmission of pathogenic agent to humans. The infection with *T. gondii* causes abortion and congenital malformations in pregnant humans (El-Sherbini et al., 2019).

#### **Fungal zoonoses**

Various fungal infections are known in humans. Some fungi are air born and the environment has an important role in their distribution. A number of fungal pathogens have zoonotic potential and public health importance transmitting naturally between birds, animals, and humans (Gnat et al., 2021). Infectious fungi can affect humans as a primary pathogen or opportunists. In most cases, fungal zoonoses develop no or mild clinical signs. However, immunosuppressive patients show severe diseases that can lead to death (Seyedmousavi, 2015). *Aspergillus* spp. and *Candida* spp. transmit from pet birds to humans

that cause severe diseases in immunocompromised patients (Cray, 2011; Boseret et al., 2013).

Among the zoonotic diseases, fungal zoonoses have not received enough attention at the international public health level, leading to insufficient attention to their preventive strategies (Seyedmousavi, 2015). Infected birds contaminate the soil and the environment, and fungal pathogens are mainly transmitted to human by direct or indirect contact with sick birds, exposure to contaminated environment, and inhalation of aerosols and respiratory particles (Richard et al., 2017). Some fungi like *Histoplasmosis* spp. and *Cryptococcus* spp. transfer from birds to soil and grow in soil, and consequently the contaminated soil can be a way for zoonoses transmission (Rahman et al., 2020). The most common fungal zoonoses from birds are dermatophytosis, histoplasmosis and cryptococcosis, which are explained below.

#### ***Cryptococcosis***

Cryptococcosis is an infectious zoonotic disease by *Cryptococcus* spp. (Lugarini, 2008). *C. neoformans* is isolated from bird excreta, and can transmit to humans. Several birds, including chickens, pigeons, and pheasants, are hosts for *Cryptococcus* spp. (Gnat et al., 2021). Pigeons and parrots (mainly different cockatoo species) are the primary hosts of *C. neoformans* (Lugarini, 2008). Also, other pet birds, such as canaries, lovebirds, budgerigars, and cockatiels, are susceptible to *C. neoformans* as potential reservoirs for the fungus (Brilhante et al., 2010). Mostly, sick birds have no significant clinical signs except increasing body temperature and fever. The cutaneous lesions has been reported in cockatoos infected by *C. neoformans*. Birds shed the fungus through feces, and contaminated cages and soil with birds' feces are transmission routes for humans. Infection in people can occur through the inhalation of infected particles (Hashemi et al., 2014).

Infection by *C. neoformans* in humans is commonly an opportunistic infection. Meningoencephalitis often occurs in immunosuppressed persons (HIV/AIDS patients and organ transplant recipients). The infected

patients may show clinical symptoms such as headaches, fever, malaise, stiff neck, nervous signs, and even death. Respiratory and cutaneous forms of cryptococcosis can also occur in infected humans (Pasa et al., 2012; Hayashida et al., 2017; Zhang et al., 2019). Cutaneous nodular lesions have been reported in immunosuppressed pet cockatoo owner by infection with *C. neoformans* (Nosanchuk et al., 2000).

#### ***Histoplasmosis***

Histoplasmosis is a fungal disease caused by *Histoplasma capsulatum*, which is isolated from feces of various animals and birds. Pigeons and doves are the most common hosts for *H. capsulatum*. The bat droppings, feces of infected birds, and birds' nests (especially starlings) are the sources of infection for humans (Haag-Wackernage et al., 2004; Adebisi and Oluwayelu, 2018). This fungus can be persisted in soil and suitable environments for several years and transmitted to humans through contaminated soil (Rahman et al., 2020). Also, inhalation of fungal spores is another route of transmission to humans. Incubation period in human infections is about 3-17 days. Mostly, *H. Capsulatum* infection in human occurs as an asymptomatic disease. In a few cases, respiratory symptoms including malaise, headache, fever, dry cough, and chest pain can develop. Pneumonia occurs in rare cases, usually seen in untreated infected people that may even lead to death. In some patients, it develops skin ulcers and lymphadenopathy (Nagoda et al., 2012; Adebisi and Oluwayelu, 2018).

#### ***Dermatophytosis***

Dermatophytes, known as ringworm fungi, are the most common isolated fungi from birds and animals (Nweze, 2010). Dermatophytes have three different genera, including *Trichophyton*, *Epidermophyton*, and *Microsporum* spp. (Popoola et al., 2006; Nweze, 2010). *Trichophyton* and *Microsporum* spp. are zoophilic, causing skin lesions in animals and birds. Contact with infected birds, feathers, and skin debris represents a potential role of fungi transmission to humans. In people with dermatophytosis, skin lesions can be seen as red areas, along with skin itching and

central pallor, which appears as a "ring" (Fehr, 2015; Adebiyi and Oluwayelu, 2018).

### Prevention and control of transmission from birds to human

The occurrence of zoonotic diseases depends on the interaction of birds, humans, and the environment. Therefore, managing and performing effective protocols are required for the control and prevention of these diseases (Chomel, 2009). Restricting human contact with sick birds and hand washing with disinfectant sprays and gels are the major crucial strategies to control zoonoses (Al-Tayib et al., 2019). Identifying, quarantining, and treating sick birds help to prevent zoonoses. Cleaning and disinfecting the birds' cage, food and water bowls with a bleach solution decreases the risk of infection and transmission of zoonotic pathogens to humans. Keeping pet birds away from high-risk individuals (such as very young and very old people or patients with weakened immune systems) will help reduce their risk of infection (Aenishaenslin et al., 2013; Dahal and Kahn, 2014). Detection and eradication of carriers, biological and mechanical vectors can avoid the transmission of zoonotic agents between birds and humans (Boseret et al., 2013; Rahman et al., 2020).

### Conclusion

Zoonotic diseases associated with pet birds do not mainly lead to a severe infection in healthy humans, but immunocompromised individuals are at serious risk for infection by zoonotic pathogens. Exposure to infected pet birds or their secretions and droppings leads to transmitting zoonoses to humans. Identifying and treating sick birds and applying hygiene protocols significantly decrease zoonotic infection risk in people.

### Acknowledgments

Not applicable.

### Ethical approval

Not applicable.

### Conflicts of interest

The author declares no conflict of interest.

### References

- Adebiyi A. I. & Oluwayelu D. O. Zoonotic fungal diseases and animal ownership in Nigeria. *Alexandria Journal of Medicine*, 2018, 54(4), 397-402.
- Aenishaenslin C., Hongoh V., Cissé H. D., Hoen A. G., Samoura K., Michel P., Waaub J. P. & Bélanger D. Multi-criteria decision analysis as an innovative approach to managing zoonoses: Results from a study on Lyme disease in Canada. *BMC Public Health*, 2013, 13, 897.
- Alazawy A. K. & Al Ajeeli K. S. Isolation and molecular identification of wild Newcastle disease virus isolated from broiler farms of Diyala Province, Iraq. *Veterinary World*, 2020, 13(1), 33-9.
- Al-Tayib O.A. An overview of the most significant zoonotic viral pathogens transmitted from animal to human in Saudi Arabia. *Pathogens*, 2019, 8, 25.
- Ammar S., Wood L., Su C., Spriggs M., Brown J., Van Why K. & Gerhold R. Toxoplasma gondii prevalence in carnivorous wild birds in the eastern United States. *International Journal for Parasitology: Parasites and Wildlife*, 2021, 15, 153-7.
- Anamaria M. P., Rafaela S., Junior C. A. C. Virulence factors in *Salmonella typhimurium*: the sagacity of a bacterium. *Current Microbiology*, 2018, 76(6), 762-73.
- Andino A. & Hanning I. *Salmonella enterica*: survival, colonization, and virulence differences among Serovars. *Science World Journal*, 2015, 520179.
- Antilles N., Sanglas A. & Cerdà-Cuellar M. Free-living waterfowl as a source of zoonotic bacteria in a dense wild bird population area in northeastern Spain. *Transboundary and Emerging Diseases*, 2015, 62, 516–21.
- Antillón M., Warren J. L., Crawford F. W., Weinberger D. M., Kürüm E., Pak G. D., Pak G. D., Marks F. & Pitzer V. E. The burden of typhoid fever in low- and middle-income countries: A meta-regression approach. *PLoS Neglected Tropical Diseases*, 2017, 11, 1-21.
- Antunes P., Mourão J., Campos J. & Peixe L. Salmonellosis: the role of poultry meat. *Clinical Microbiology and Infection*, 2016, 22, 110-21.
- Awofisayo-Okuyelu A., Hall I., Adak G., Hawker J., Abbott S. & McCarthy N. A systematic review and meta-analysis on the incubation

- period of Campylobacteriosis. *Epidemiology and Infection*, 2017, 145(11), 2241-53.
- Backert S., Tegtmeyer N., Cróinín T.Ó., Boehm M., Heimesaat M. M. Human campylobacteriosis. In: Klein G (ed.) *Campylobacter*. Academic, 2017, 1-25.
- Bai F. E. A., Thompson P. J. S. & Leis A. A. Current understanding of west Nile virus clinical manifestations, immune responses, neuroinvasion, and immunotherapeutic implications. *Pathogens*, 2019, 8, 193.
- Barbour E. K., Ayyash D. B., Alturkistni W., Alyahiby A., Yaghmoor S., Iyer A., Yousef J., Kumosani T. & Harakeh S. Impact of sporadic reporting of poultry *Salmonella* serovars from selected developing countries. *Journal of Infection in Developing Countries*, 2015, 9(1), 1-7.
- Barstad B. Direct molecular detection and genotyping of *Borrelia burgdorferi sensu lato* in the cerebrospinal fluid of children with Lyme neuroborreliosis. *Journal of Clinical Microbiology*, 2018, 56, e01868-17.
- Behboudi E. & Hamidi Sofiani V. Immune responses to Newcastle disease virus as a minor zoonotic viral agent. *Journal of Zoonotic Diseases*, 2021, 5(4), 12-23.
- Belser J. A., Lash R. R., Garg S., Tumpey T. M. & Maines T. R. The eyes have it: Influenza virus infection beyond the respiratory tract. *The Lancet Infectious Diseases*, 2018, 18, e220-e227.
- Bi Y., Chen Q., Wang Q., Chen J., Jin T., Wong G., Quan C., Liu J., Wu J., Yin R., Zhao L., Li M., Ding Z., Zou R., Xu W., Li H., Wang H., Tian K. & Gao G. F. Genesis, evolution and prevalence of h5n6 avian influenza viruses in China. *Cell Host and Microbe*, 2016, 20(6), 810-21.
- Boroomand Z. & Faryabi. S. Bird zoonotic diseases. *Journal of Zoonotic Diseases*, 2020, 4(3), 20-33.
- Boseret G., Losson B., Mainil J. G., Thiry E. & Saegerman C. Zoonoses in pet birds: review and perspectives. *Veterinary Research*, 2013, 44, 36.
- Braima K., Zahedi A., Oskam C., Reid S., Pingault N., Xiao L. & Ryan U. Retrospective analysis of *cryptosporidium* species in western Australian human populations (2015-2018), and emergence of the *C. hominis* IfA12G1R5 subtype. *Infection, Genetics and Evolution*, 2019, 73, 306-13.
- Brilhante R. S. N., Castelo-Branco D. S. C. M., Soares G. D. P., Astete-Medrano D. J., Monteiro A. J., Cordeiro R. A., Sidrim J. J. C. & Rocha M. F. G. Characterization of the gastrointestinal yeast microbiota of cockatiels (*Nymphicus hollandicus*): a potential hazard to human health. *Journal of Medical Microbiology*, 2010, 59: 718-23.
- Brooks P. T., Brakel K. A., Bell J. A., Bejcek C. E., Gilpin T., Brudvig J. M. & Mansfield L. S. Transplanted human fecal microbiota enhanced Guillain Barre syndrome autoantibody responses after *Campylobacter jejuni* infection in C57BL/6 mice. *Microbiome*, 2017, 5 (1), 92.
- Bryan E. R., McLachlan R. I., Rombauts L., Katz D. J., Yazdani A., Bogoevski K., Chang C., Giles M. L., Carey A. J., Armitage C. W, Trim L. K, McLaughlin E. A. & Beagley K. W. Detection of *chlamydia* infection within human testicular biopsies. *Human Reproduction*, 2019, 34(10), 1891-8.
- Cano L., de Lucio A., Bailo B., Cardona G. A., Muadica A. S., Lobo L. & Carmena D. Identification and genotyping of *Giardia* spp. and *Cryptosporidium* spp. isolates in aquatic birds in the Salburua wetlands, Alava, Northern Spain. *Veterinary Parasitology*, 2016, 221, 144-8.
- Chancey C., Grinev A., Volkova E. & Rios M. The global ecology and epidemiology of West Nile virus. *BioMed Research International*, 2015, 376230.
- Chang C. C., Wechtaisong W., Chen S. Y., Cheng M. C., Chung C. S., Lin L. S., Lien, Y. Y. & Tsai Y. L. Prevalence and risk factors of zoonotic dermatophyte infection in pet rabbits in northern Taiwan. *Journal of Fungi*, 2022, 8, 627.
- Chen H., Yuan H., Gao R., Zhang J., Wang D., Xiong Y., Fan G., Yang F., Li X., Zhou J., Zou S., Yang L., Chen T., Dong L., Bo H., Zhao X., Zhang Y., Lan Y., Bai T., Dong J., Li Q., Wang S., Zhang Y., Li H., Gong T., Shi Y., Ni X., Li J., Zhou J., Fan J., Wu J., Zhou X., Hu M., Wan J., Yang W., Li D., Wu G., Feng Z., Gao G.F., Wang Y., Jin Q., Liu M. & Shu Y. Clinical and epidemiological characteristics of a fatal case of avian influenza A H10N8 virus infection: A descriptive study. *Lancet*, 2014, 383, 714-21.
- Cheong H. C., Lee C. Y. Q., Cheok Y. Y., Tan G. M. Y., Looi C. Y. & Wong W. F.

- Chlamydiaceae: Diseases in Primary Hosts and Zoonosis. *Microorganisms*, 2019, 7, 146.
- Chomel B. B. & Sun B. Zoonoses in the bedroom. *Emerging Infectious Diseases*, 2001, 17,167-72.
- Chomel B. B. 2009, Zoonoses. In: *Encyclopedia of Microbiology*, C. A. Davis. 3rd ed. Elsevier Inc., University of California, USA, 820-9.
- Circella E., Pugliese N., Todisco G., Cafiero M. A., Sparagano O. A. E. & Camarda A. *Chlamydia psittaci* infection in canaries heavily infested by *Dermanyssus gallinae*. *Experimental and Applied Acarology*, 2011, 55, 329-38.
- Cohen E. B., Auckland L. D., Marra P. P., Hamer S. A. Avian migrants facilitate invasions of Neotropical ticks and tick-borne pathogens into the United States. *Applied and Environmental Microbiology*, 2015, 81(24), 8366-78.
- Coipan E. C., Jahfari S., Fonville M., Oei G. A., Spanjaard L., Takumi K., Hovius J. W. R. & Sprong H. Imbalanced presence of *Borrelia burgdorferi* s.l. *multilocus* sequence types in clinical manifestations of Lyme borreliosis. *Infection, Genetics and Evolution*, 2016, 42, 66-76.
- Cong W., Huang S. Y., Zhou D. H., Xu M. J., Wu S. M., Yan C., Zhao Q., Song H. & Zhu X. First report of *Toxoplasma gondii* infection in market-sold adult chickens, ducks and pigeons in Northwest China. *Parasites & Vectors*, 2012, 5:110.
- Contreras A., Gómez-Martín A., Paterna A., Tatay-Dualde J., Prats-van der Ham M., Corrales J.C., de la Fe C. & Sánchez A. Epidemiological role of birds in the transmission and maintenance of zoonoses. *Revue Scientifique et Technique*, 2016, 35, 3.
- Cray C. Infectious and zoonotic disease testing in pet birds. *Clinical and Laboratory Medicine Journal*, 2011, 31, 71-85.
- Da Cunha M. J. R., Cury M. C. & Santin M. Molecular identification of *Enterocytozoon bieneusi*, *Cryptosporidium*, and *Giardia* in Brazilian captive birds. *Parasitology Research*, 2017, 116, 487-93.
- Dahal R. & Kahn L. Zoonotic diseases and one health approach. *Epidemiology*, 2014, 10.
- David S. & Abraham A. M. Epidemiological and clinical aspects on West Nile virus, a globally emerging pathogen. *Infectious Diseases*, 2016, 48, 571-86.
- Day M. J. Pet-Related Infections. *American Family Physician*, 2016, 94, 794-802.
- Dhama K., Mahendran M., Tiwari R., Singh S. D., Kumar D., Singh S. & Sawant P. M. Tuberculosis in Birds: Insights into the *Mycobacterium avium* Infections. *Veterinary Medicine International*, 2011, 712369.
- Dorrestein G. M. Bacterial and parasitic diseases of passerines. *Veterinary Clinics of North America: Exotic Animal Practice*, 2009, 12, 433-51.
- Dumas A., Bouchard C., Dibernardo A., Drapeau P., Lindsay L.R., Ogden N. H. & Leighton P. A. Transmission patterns of tick-borne pathogens among birds and rodents in a forested park in southeastern Canada. *PLoS One*, 2022, 17(4), e0266527.
- Ebani V. V., Guardone L., Bertelloni F., Perrucci S., Poli A. & Mancianti F. Survey on the presence of bacterial and parasitic zoonotic agents in the feces of wild birds. *Veterinary Sciences*, 2021, 8, 171.
- Ehuwa O., Jaiswal A. K. & Jaiswal S. *Salmonella*, food safety and food handling practices. *Foods*, 2021, 10, 907.
- El-Sherbini M. S., Abd El-Aal A. A., El-Sherbiny W. S., Attia S. S., Abdel Aziz I. Z., Nasr G. M., Salama M. S. & Badr M. S. Toxoplasmosis and abortion: pro- and anti-inflammatory cytokines gene expression of the host immune cells. *Egyptian Journal of Medical Human Genetics*, 2019, 20, 3.
- El-Tayeb M., Ibrahim A. S. S., Al-Salamah A. A., Almaary K. S. & Elbadawi Y.B. Prevalence, serotyping and antimicrobials resistance mechanism of *Salmonella enterica* isolated from clinical and environmental samples in Saudi Arabia. *Brazilian Journal of Microbiology*, 2017, 48, 499-508.
- Fehr M. Zoonotic potential of dermatophytosis in small mammals. *Journal of Exotic Pet Medicine*, 2015, 24 (3), 308-16.
- Fernandez-Cruz A., Munoz P., Mohedano R., Valerio M., Marín M., Alcalá L., Rodriguez-Créixems M., Cercenado E. & Bouza E. *Campylobacter* bacteremia: clinical characteristics, incidence, and outcome over 23 years. *Medicine*, 2010, 89, 319-30.
- França M. S. and Brown J. D. 2014, *Influenza pathobiology and pathogenesis in avian species*. Springer International Publishing, New York, NY.

- French N. P., Midwinter A., Holland B., Collins-Emerson J., Pattison R., Colles F. & Carter P. Molecular epidemiology of *Campylobacter jejuni* Isolates from wild-bird fecal material in children's playgrounds. *Applied and Environmental Microbiology*, 2009, 75, 779.
- Fulton R. M. & Sanchez S. 2018, Other Bacterial Diseases. In: *Diseases of Poultry*. Swayne D. E., Boulianne M., Logue C. M., McDougald L. R., Nair V., Suare D. L., Eds. 14th ed. American Association of Avian Pathologists: Hoboken, NJ, USA, 1033-1043.
- Gaukler S. M., Linz G. M., Sherwood J. S., Dyer N. W., Bleier W. J., Wannemuehler Y. M., Nolan L. K. & Logue C. M. *Escherichia coli*, *salmonella*, and *Mycobacterium avium* subsp. *paratuberculosis* in wild European starlings at a Kansas cattle feedlot. *Avian Disease*, 2009, 53, 544-51.
- Giotis E. S., Carnell G., Young E. F., Ghanny S., Soteropoulos P., Wang L. F., Barclay W. S., Skinner M. A. & Temperton N. Entry of the bat influenza H17N10 virus into mammalian cells is enabled by the MHC class II HLA-DR receptor. *Nature Microbiology*, 2019, 4, 2035-38.
- Gnat S., Łagowski D., Nowakiewicz A. & Dylağ M. A global view on fungal infections in humans and animals: opportunistic infections and microsporidiosis. *Journal of Applied Microbiology*, 2021, 131 (5), 2095-113.
- Gow A. G., Gow D. J., Hall E. J., Langton D., Clarke C. & Pappasouliotis K. Prevalence of potentially pathogenic enteric organisms in clinically healthy kittens in the UK. *Journal of Feline Medicine and Surgery*, 2009, 655-62.
- Gray G. C., Ferguson D. D., Lowther P. E., Heil G. L. & Friary J. A. A national study of us bird banders for evidence of avian influenza virus infections. *Journal of Clinical Virology*, 2011, 51, 132-5.
- Guo Y., Ryan U., Feng Y. & Xiao L. Association of common zoonotic pathogens with concentrated animal feeding operations. *Frontiers in Microbiology*, 2022.
- Haag-Wackernage D. & Moch H. Health hazards posed by feral pigeons. *The Journal of Infection*, 2004, 48(4), 307-13.
- Hamer S. A., Goldberg T. L., Kitron U. D., Brawn J. D., Anderson T. K., Loss S. R., Walker E. D. & Hamer G. L. Wild birds and urban ecology of ticks and tick-borne pathogens, Chicago, Illinois, USA, 2005–2010. *Emerging Infectious Diseases*, 2012, 18(10), 1589-95.
- Hao Q., Hou X., Geng Z. & Wan K. Distribution of *Borrelia burgdorferi sensu lato* in China. *Journal of Clinical Microbiology*, 2011, 49(2), 647-50.
- Haridy M., Fukuta M., Mori Y., Ito H., Kubo M., Sakai H. & Yanai T. An Outbreak of Mycobacterium genavense Infection in a Flock of Captive Diamond Doves (*Geopelia cuneata*). *Avian Disease*, 2014, 58, 383-90.
- Hashemi S. J., Jabbari A. G., Bayat M. & Mashhadi Rafiei S. Prevalence of *Cryptococcus neoformans* in domestic birds referred to veterinary clinics in Tehran. *European Journal of Experimental Biology*, 2014, 4(1), 482-6.
- Hawkins M. G., Crossley, B. M., Osofsky A., Webby R.J., Lee C.W., Suarez D.L. & Hietala S. K. Avian influenza A virus subtype H5N2 in a red-lored Amazon parrot. *Journal of the American Veterinary Medical Association*, 2006, 228 (2), 236-4.
- Hayashida M. Z., Seque C. A., Pasin V. P., Enokihara M. M. S. & Porro A. M. Disseminated cryptococcosis with skin lesions: report of a case series. *Anais Brasileiros de Dermatologia*, 2017, 92, 69-72.
- Heesterbeek H., Anderson R. M. & Andreasen V. Modeling infectious disease dynamics in the complex landscape of global health. *Science*, 2015, 347 (4339), aaa4339.
- Hill S. C., Hansen R., Watson S., Coward V., Russell C., Cooper J., Essen S., Everest H., Parag K. V., Fiddaman S., Reid S., Lewis N., Brookes S. M., Smith A. L., Sheldon B., Perrins C. M., Brown I. H. & Pybus O.G. Comparative micro-epidemiology of pathogenic avian influenza virus outbreaks in a wild bird population. *Philosophical Transactions of the Royal Society B*, 2019, 374:20180259.
- Hogerwerf L., De Gier B., Baan B. & Van Der Hoek W. *Chlamydia psittaci* (psittacosis) as a cause of community-acquired pneumonia: a systematic review and meta-analysis. *Epidemiology and Infection*, 2017, 145(15), 3096-105.
- Hollingsworth T. D., Pulliam J. R. C., Funk S., Truscott J. E., Isham V. & Lloyd A. L. Seven challenges for modelling indirect transmission: vector-borne diseases,



- macroparasites and neglected tropical diseases. *Epidemics*, 2015, 10, 16-20.
- Hoop R. K. Mycobacterium tuberculosis infection in a canary (*Serinus canana* L.) and a blue-fronted Amazon parrot (*Amazona amazona aestiva*). *Avian Disease*, 2002, 46(2), 502-4.
- Hsieh Y. H. & Sulaiman I. M. Campylobacteriosis: An emerging infectious foodborne disease. *Foodborne Disease*, 2018, 119-55.
- Huang S. Y., Fan Y. M., Chen K., Yao Q.X. & Yang B. Seroprevalence and risk assessment of *Toxoplasma gondii* in Java sparrows (*Lonchura oryzivora*) in China. *BMC Veterinary Research*, 2019, 15, 129.
- Hung Y. T., Lay C. J., Wang C. L. & Koo M. Characteristics of nontyphoidal gastroenteritis in Taiwanese children: A 9-year period retrospective medical record review. *Journal of Infection and Public Health*, 2017, 10, 518-21.
- Iannino F., Salucci S., Di Donato G., Badagliacca P., Vincifori G., Di Giannatale E. *Campylobacter* and antimicrobial resistance in dogs and humans: "one health" in practice. *Veterinaria Italiana*, 2019, 55(3), 203–20.
- Jacob J. & Lorber B. Diseases transmitted by man's best friend: The dog. *Microbiology Spectrum*, 2015, 3, 111-31.
- Jajere S. M. A review of *Salmonella enterica* with particular focus on the pathogenicity and virulence factors, host specificity and adaptation and antimicrobial resistance including multidrug resistance. *Veterinary World*, 2019, 12(4), 504-21.
- Jarvis N. A., O'Bryan C. A., Dawuod T. M., Hong Park S., Min Kwon Y., Crandall P. G. & Ricke S. C. An overview of *Salmonella* thermal destruction during food processing and preparation. *Food Control*, 2016, 68, 280-90.
- Jiao P., Song Y., Yuan R., Wei L., Cao L., Luo K. & Liao M. Complete genomic sequence of an H5N1 influenza virus from a parrot in southern China. *Journal of Virology*, 2012a, 86 (16), 8894-95.
- Jiao P., Wei L., Yuan R., Gong L., Cao L., Song Y., Luo K., Ren T. & Liao M. Complete genome sequence of an H5N2 avian influenza virus isolated from a parrot in southern China. *Journal of Virology*, 2012b, 86 (16), 8890-91.
- Johansson H., Ellström P., Artursson K., Berg C., Bonnedahl J., Hansson I., Hernandez J., Lopez-Martín J., Medina-Vogel G., Moreno L., Olsen B., Olsson Engvall E., Skarin H., Troell K., Waldenström J., Ågren J. & González-Acuña D. Characterization of *Campylobacter* spp. isolated from wild birds in the Antarctic and Sub-Antarctic. *PLoS One*, 2018, 13:e0206502.
- Jones F. T. A review of practical Salmonella control measures in animal feed. *Journal of Applied Poultry Research*, 2011, 20, 102-13
- Jorn K. S., Thompson K. M., Larson J. M. & Blair J. E. Polly can make you sick: Pet bird-associated diseases. *Cleveland Clinic Journal of Medicine*, 2009, 76, 235-43.
- Kaakoush N. O., Castaño-Rodríguez N., Mitchell H. M. & Man S. M. Global epidemiology of *Campylobacter* infection. *Clinical Microbiology Reviews*, 2015, 28, 687-720.
- Kallapura G., Morgan M. J., Pumford N. R., Bielke L. R., Wolfenden A. D., Faulkner O. B., Latorre J. D., Menconi A., Hernandez-Velasco X., Kuttappan V.A., Hargis B. M. & Tellez G. Evaluation of the respiratory route as a viable portal of entry for *Salmonella* in poultry via intratracheal challenge of *Salmonella Enteritidis* and *Salmonella Typhimurium*. *Poultry Science*, 2014, 93, 340-6.
- Kampen H., Holicki C. M., Ziegler U., Groschup M. H., Tews B. A. & Werner D. West Nile Virus Mosquito Vectors in Germany. *Viruses*, 2020, 12, 493.
- Kauffman M. D. & LeJeune J. European starlings (*Sturnus vulgaris*) challenged with *Escherichia coli* O157 can carry and transmit the human pathogen to cattle. *Letters in Applied Microbiology*, 2011, 53, 596-601.
- Karimi Madab M., Dadras H., Hosseini A., Hosseinian S. A. & Sepehrimanesh M. Partial cloning and expression of HA1 and HA2 gene of avian influenza H9N2 gene in *Lactococcus lactis*. *Online Journal Of Veterinary Research*, 2013, 261-9.
- Kirk M. D., Pires S. M., Black R. E., Caipo M., Crump J. A., Devleeschauwer B., Döpfer D., Fazil A., Fischer-Walker C. L., Hald T., Hall A. J., Keddy K. H., Lake R. J., Lanata C. F., Torgerson P. R., Havelaar A. H. & Angulo F. J. World Health Organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: a data synthesis. *PLoS Medicine*, 2015, 12, e1001921.

- Kozdruń W., Czekaj H. & Styś N. Avian zoonoses - a review. *Bulletin of the Veterinary Institute in Pulawy archives*, 2015, 59, 171-8.
- Kuiken T., Breitbart M., Beer M., Grund C., Höper D., Van den Hoogen B., Kerkhoffs J. H., Kroes A. C. M., Rosario K., van Run P., Schwarz M., Svraka S., Teifke J. & Koopmans M. Zoonotic Infection With Pigeon Paramyxovirus Type 1 Linked to Fatal Pneumonia. *The Journal of Infectious Diseases*, 2018, 218 (7), 1037-44.
- Kürekci C., Sakin F., Epping L., Knüver M.T., Semmler T. & Stingl K. Characterization of *Campylobacter* spp. Strains Isolated From Wild Birds in Turkey. *Frontiers in Microbiology*, 2021, 12, 712106.
- Kurtz J. R., Goggins J. A. & McLachlan J. B. *Salmonella* infection: Interplay between the bacteria and host immune system. *Immunology Letters*, 2017, 190, 42-50
- Lai S., Qin Y., Cowling B. J., Ren X., Wardrop N. A., Gilbert M., Tsang T. K., Wu P., Feng L., Jiang H. Peng Z., Zheng J., Liao Q., Li S., Horby P. W., Farrar J. J., Gao G. F., Tatem A. J. & Yu H. Global epidemiology of avian influenza A H5N1 virus infection in humans, 1997-2015: A systematic review of individual case data. *The Lancet Infectious Diseases*, 2016, 16, e108-e118.
- Lang A. S., Lebarbenchon C., Ramey A. M., Robertson G. J., Waldenström J. & Wille M. Assessing the role of seabirds in the ecology of influenza A viruses. *Avian Disease*, 2016, 60(1s), 378-86.
- Lawson B., De Pinna E., Horton R. A., MacGregor S. K., John S. K., Chantrey J., Duff J. P., Kirkwood J. K., Simpson V. R., Robinson R. A., Wain J. & Cunningham A. A. Epidemiological evidence that garden birds are a source of human salmonellosis in England and Wales. *PLoS One*, 2014, 9, e88968.
- Lawson B., De Pinna E., Horton R. A., Macgregor S. K. & John S. K. Epidemiological evidence that garden birds are a source of human salmonellosis in England and wales. *PLoS One*, 2014, 9(2), e88968.
- Lennox A. M. Mycobacteriosis in Companion Psittacine Birds: A Review. *Journal of Avian Medicine and Surgery*, 2007, 21(3), 181-7.
- Li X., Zhang R., Huang Z., Yao D., Luo L., Chen J., Ye W., Li L., Xiao S., Liu X., Ou X., Sun B., Xu M., Yang R. & Zhang X. Estimation of avian influenza viruses in water environments of live poultry markets in Changsha, China, 2014 to 2018. *Food and Environmental Virology*, 2022, 14(1), 30-9.
- Linde A. M., Munir M., Zohari S., Stahl K., Baule C., Ståhl K., Baule C., Renström L. & Berg M. Complete genome characterization of a Newcastle disease virus isolated during an outbreak in Sweden in 1997. *Virus Genes*, 2011, 41(2), 165-73.
- Long J. S., Mistry B., Haslam S. M. & Barclay W. S. Host and viral determinants of influenza A virus species specificity. *Nature Reviews Microbiology*, 2019, 17, 67-81.
- Lugarini C., Goebel C. S., Condas L. A. Z., Muro M. D., De Farias M. R., Ferreira F. M., & Vainstein M. H. *Cryptococcus neoformans* isolated from Passerine and Psittacine bird excreta in the state of Paraná, Brazil. *Mycopathologia*, 2008, 166, 61-9.
- Maësaar M., Tedersoo T., Meremaë K. & Roasto M. The source attribution analysis revealed the prevalent role of poultry over cattle and wild birds in human campylobacteriosis cases in the Baltic States. *PLoS One*, 2020, 15:e0235841.
- Magouras I., Brookes V. J., Jori F., Martin A., Pfeiffer D. U. & Dürr S. Emerging Zoonotic Diseases: Should We Rethink the Animal–Human Interface? *Frontiers in Veterinary Science*, 2020, 7.
- Malik Y. S., Arun Prince Milton A., Ghatak S. & Ghosh S. Avian Chlamydiosis (Psittacosis, Ornithosis). In: *Role of birds in transmitting zoonotic pathogens. Livestock Diseases and Management*, 2021
- Marzal A., Ferraguti M., Muriel J., Magallanes S., Ortize J. A., García-Longoria L., Bravo-Barriga D., Guerrero-Carvajal F., Aguilera-Sepúlveda P., Llorente F., De Lope F., Jiménez-Clavero M. Á. & Frontera E. Circulation of zoonotic flaviviruses in wild passerine birds in Western Spain. *Veterinary Microbiology*, 2022, 268, 109399.
- Michel F., Fischer D., Eiden M., Fast C., Reuschel M., Müller K., Rinder M., Urbaniak S., Brandes F., Schwehn R., Lühken R., Groschup M. H. & Ziegler U. West Nile Virus and Usutu Virus Monitoring of Wild Birds in Germany. *International Journal of Environmental Research and Public Health*, 2018, 15, 171.

- Modirihamedan A., Aghajantabar S., King J., Graaf A., Pohlmann A., Aghaiyan L., Ziafati Z., Mahfoozi Y., Hosseini H., Beer M., Ghalyanchi langeroudi A. & Harder T. Wild bird trade at live poultry markets potentiates risks of avian influenza virus introduction in Iran, *Infection Ecology & Epidemiology*, 2021, 11, 1.
- Mohan V., Stevenson M., Marshall J., Fearnhead P., Holland B.R., Hotter G. & French N. P. *Campylobacter jejuni* colonization and population structure in urban populations of ducks and starlings in New Zealand. *Microbiologyopen*, 2013, 2, 659-73.
- Mostafa A., Abdelwhab E. M., Mettenleiter T. C. & Pleschka S. Zoonotic potential of influenza A viruses: a comprehensive overview. *Viruses*, 2018, 10, 497.
- Mughini-Gras L., Pijnacker R., Coipan C., Mulder A. C., Veludo A. F., De Rijk S., Van Hoek A. H. A. M., Buij R., Muskens G., Koene M., Veldman K., Duim B., Van der Graaf-van Bloois L., Van der Weijden C., Kuiling S., Verbruggen A., Van der Giessen J., Opsteegh M., Van der Voort M., Castelijin G. A. A., Schets F. M., Blaak H., Wagenaar J. A., Zomer A. L. & Franz E. Sources and transmission routes of campylobacteriosis: a combined analysis of genome and exposure data. *The Journal of Infectious Diseases*, 2021, 82, 216-26.
- Mulder A. C., Franz E., De Rijk S., Versluis M. A. J., Coipan C., Buij R., Muskens G., Koene M., Pijnacker R., Duim B., Van der Graaf-van Bloois L., Veldman K., A Wagenaar J., Zomer A. L., Schets F. M., Blaak H. & Mughini-Gras L. Tracing the animal sources of surface water contamination with *Campylobacter jejuni* and *Campylobacter coli*. *Water Research*, 2020, 187, 116421.
- Munir S., Hussain M., Farooq U., Ullah Z., Jamal Q., Afreen M., Bano K., Khan J., Ayaz S., Yong Kim K. & Anees M. Quantification of antibodies against poultry haemagglutinating viruses by haemagglutination inhibition test in Lahore. *African Journal of Microbiology Research*, 2012, 6(21), 4614-19.
- Mygland Å., Løstad U., Fingerle V., Rupprecht T., Schmutzhard E. & Steiner I. EFNS guidelines on the diagnosis and management of European Lyme neuroborreliosis. *European Journal of Neurology*, 2009, 17, 8-e4.
- Nagoda M., Uloko A.E., Babashani M. & Maiyaki M. Histoplasmosis: An elusive re-emerging chest infection. *Nigerian Journal of Clinical Practice*, 2012, 15 (2), 235-7.
- Narayanan M. S., Parthiban M., Sathiya P. & Kumanan K. Molecular detection of Newcastle disease virus using flinders molecular detection of Newcastle disease virus using flinders technology associates-PCR technology associates-PCR. *Veterinarski Arhiv*, 2010, 80(1), 51-60.
- Nidaullah H., Abirami N., Shamila-Syuhada A. K., Chuah L. O., Nurul H., Tan, T. P., Abidin F. W. Z. & Rusul G. Prevalence of *Salmonella* in poultry processing environments in wet markets in Penang and Perlis, Malaysia. *Veterinary World*, 2017, 10, 286-92.
- Nosanchuk J. D., Shoham S., Fries B., Shapiro D. & Levitz S.M. Evidence of zoonotic transmission of *Cryptococcus neoformans* from a pet cockatoo to an immunocompromised patient. *Annals of Internal Medicine*, 2000, 132(3), 205-8.
- Nweze E. I. Dermatophytosis in Western Africa: a review. *Pakistan Journal of Biological Sciences: PJBS*, 2010, 13(13), 649-56.
- Ogden N., Barker I., Francis C., Heagy A., Lindsay L. & Hobson K. How far north are migrant birds transporting the tick *Ixodes scapularis* in Canada? Insights from stable hydrogen isotope analyses of feathers. *Ticks and Tick-Borne Diseases*, 2015, 6(6), 715-20.
- Osek J. & Wiczorek K. Zoonoses and zoonotic agents in Europe in 2012—Report of the European Food Safety Authority (EFSA). *Życie Weterynaryjne*, 2014, 89, 472-8.
- Pan X., Su X., Ding P., Zhao J., Cui H., Yan D., Teng Q., Li X., Beerens N., Zhang H., Liu Q., de Jong M.C.M. & Li Z. Maternal-derived antibodies hinder the antibody response to H9N2 AIV inactivated vaccine in the field. *Animal Diseases*, 2022.
- Pasa C. R., Chang M. R. & Hans-Filho G. Post-trauma primary cutaneous cryptococcosis in an immunocompetent host by *Cryptococcus gattii* VGII. *Mycoses*, 2012, 55, e1-e3.
- Peng S. & Broom D. M. The Sustainability of Keeping Birds as Pets: Should Any Be Kept? *Animals (Basel)*, 2021, 11(2), 582.
- Perez-Ramirez E., Llorente F. & Jimenez-Clavero M. A. Experimental infections of wild birds with West Nile virus. *Viruses*, 2014, 6, 752-81.

- Pfeffer M. & Dobler G. Emergence of zoonotic arboviruses by animal trade and migration. *Parasites & Vectors*, 2010, 3, 35-10.
- Popoola T. O. S., Ojo D. A. & Alabi, R. O. Prevalence of dermatophytosis in junior secondary schoolchildren in Ogun State, Nigeria. *Mycoses*, 2006, 49(6), 499-503.
- Pui C. F., Wong W. C., Chai L. C., Tunung R., Jeyaletchumi P., Noor Hidayah M. S., Ubong A., Farinazleen M. G., Cheah Y. K. & Son R. *Salmonella*: A foodborne pathogen. *International Food Research Journal*, 2011, 18(2), 465-73.
- Rahman M. T., Sobur M. A., Islam M. S., Ievy S., Hossain M. J., El Zowalaty M. E., Rahman A. M. M. T. & Ashour H. M. Zoonotic Diseases: Etiology, Impact, and Control. *Microorganisms*, 2020, 8, 1405.
- Rahmani M., Peighambari S. M., Yazdani A. & Hojjati, P. *Salmonella* infection in birds kept in parks and pet shops in Tehran, Iran. *International Journal of Veterinary Research*, 2011, 5 (3), 145-8.
- Rahmaniar R. P., Yunita M. N., Effendi M. H. & Yanestria S. M. Encoding Gene for Methicillin Resistant *Staphylococcus aureus* (MRSA) Isolated from Nasal Swab of Dogs. *Indian Journal of Veterinary Pathology*, 2020, 97(02), 37-40.
- Ramey A. M., Hill N. J., DeLiberto T. J., Gibbs S. E., Camille Hopkins M., Lang A. S., Poulson R. L., Prosser D. J., Sleeman J. M., Stallknecht D. E. & Wan X. F. Highly pathogenic avian influenza is an emerging disease threat to wild birds in North America. *The Journal of Wildlife Management*, 2022.
- Reboredo-Fernandez A., Ares-Mazas E., Caccio S.M. & Gomez-Couso H. Occurrence of *Giardia* and *Cryptosporidium* in wild birds in Galicia (Northwest Spain). *Parasitology*, 2015, 142, 917-25.
- Reemtsma H., Holicki C. M., Fast C., Bergmann F., Eiden M., Groschup M. H. & Ziegler U. Pathogenesis of West Nile Virus Lineage 2 in Domestic Geese after Experimental Infection. *Viruses*, 2022, 14, 1319.
- Ren X., Fu Y., Xu C., Feng Z., Li M., Zhang L., Zhang J. & Liao M. High resolution melting (HRM) analysis as a new tool for rapid identification of *Salmonella* enterica serovar Gallinarum biovars Pullorum and Gallinarum. *Poultry Science*, 201, 96 (5), 1088-1093.
- Richard M., Knauf S., Lawrence P., Mather A. E., Munster V. J., Müller M. A., Smith D. & Kuiken T. Factors determining human-to-human transmissibility of zoonotic pathogens via contact. *Current Opinion in Virology*, 2017, 22, 7-12.
- Ryan U. & Cacciò S. M. Zoonotic potential of *Giardia*. *International Journal for Parasitology*, 2013, 43, 943-56.
- Sachse K., Laroucau K. & Vanrompay D. Avian Chlamydiosis. *Current Clinical Microbiology Reports*, 2015, 2, 10-21.
- Sak B., Brady D., Pelikánová M., Květoňová D., Rost M., Kostka M., Pelikánová M., Tolarová V., Hůzová Z. & Kváč M. Unapparent microsporidial infection among immunocompetent humans in the Czech Republic. *The Journal of Clinical Microbiology*, 2011, 49, 1064-70.
- Samy A. & Naguib M. M. Avian respiratory coinfection and impact on avian influenza pathogenicity in domestic poultry: field and experimental findings. *Veterinary Sciences*. 2018, 5(1), 23.
- Saravanan S., Purushothaman V., Murthy T. R. G. K., Sukumar K., Srinivasan P., Gowthaman V., Balusamy M., Atterbury R., & Kuchipudi S.V. Molecular Epidemiology of Nontyphoidal *Salmonella* in Poultry and Poultry Products in India: Implications for Human Health. *Indian Journal of Medical Microbiology*, 2015, 55, 319-26.
- Scallan E., Hoekstra R. M., Angulo F. J., Tauxe R. V., Widdowson M.A., Roy S. L., Jones J. L. & Griffin P. M. Foodborne illness acquired in the United States—major pathogens. *Emerging Infectious Diseases*, 2011, 17, 7-15.
- Schurer J. M., Mosites E., Li C., Meschke S. & Rabinowitz P. Community-based surveillance of zoonotic parasites in a ‘One Health’ world: A systematic review. *One Health*, 2016, 2, 166-74.
- Scott J. D., Anderson J. F. & Durden L. A. Widespread dispersal of *Borrelia burgdorferi*-infected ticks collected from songbirds across Canada. *Journal of Parasitology Research*, 2012, 98(1), 49-59.
- Sergeev A. A., Demina O. K., Pyankov O. V., Pyankova O. G., Agafonov A. P., Kiselev S. A., Agranovski I. E., Sergeev A. A., Shikov A. N., Shishkina L.N., Safatov A. S. & Sergeev A. N. Infection of Chickens Caused by Avian Influenza Virus A/H5N1 Delivered

- by Aerosol and Other Routes. *Transboundary and Emerging Diseases*, 2013, 60, 159-65.
- Seyedmousavi S., Guillot J., Tolooe A., Verweij P. E. & De Hoog G. S. Neglected fungal zoonoses: hidden threats to man and animals. *Clinical Microbiology and Infection*, 2015, 21, 416-25.
- Shi Y., Chen J., Shi X., Hu J., Li H. X., Wang L. Y. & Wu B. A case of *chlamydia psittaci* caused severe pneumonia and meningitis diagnosed by metagenome next-generation sequencing and clinical analysis: a case report and literature review. *BMC Infectious Diseases*, 2021, 21, 621.
- Smith K. A., Campbell C. T., Murphy J., Stobierski M. G. & Tengelsen L. A. Compendium of measures to control *Chlamydophila psittaci* infection among humans (psittacosis) and pet birds (Avian Chlamydiosis), National Association of State Public Health Veterinarians (NASPHV). *Journal of Exotic Pet Medicine*, 2010, 20, 32-45.
- Socha W., Kwasnik M., Larska M., Rola J. & Rozek W. Vector-borne viral diseases as a current threat for human and animal health—one health perspective. *Journal of Clinical Medicine*, 2022, 11, 3026.
- Spackman E. 2014, A brief introduction to avian influenza virus. In: *Animal Influenza Virus*. Ed, E. Spackman. New York: Springer, 61-8.
- Stanek G., Wormser G. P., Gray J. & Strle F. Lyme borreliosis. *Lancet*, 2012, 379, 461-73.
- Steere A. C., Strle F., Wormser G. P., Hu L. T., Branda J. A., Hovius J. W. R., Li X. & Mead P. S. Lyme borreliosis. *Nature Reviews Disease Primers*, 2016, 2, 16090.
- Steinmetz H. W., Rutz C., Hoop R. K., Grest P., Bley C. R. & Hatt J. M. Possible human-avian transmission of *Mycobacterium tuberculosis* in a green-winged macaw (*Ara chloroptera*). *Avian Disease*, 2006, 50(4), 641-5.
- Stokes H. S., Berg M. L. & Bennett A. T. D. A review of chlamydial infections in wild birds. *Pathogens*, 2021, 10, 948.
- Su S., Bi Y., Wong G., Gray G. C., Gao G. F. & Li S. Epidemiology, evolution, and recent outbreaks of avian influenza virus in China. *Journal of Virology*, 2015, 89 (17), 8671-6.
- Sun Y., Bi Y., Pu J., Hu Y., Wang J., Gao H., Liu L., Xu Q., Tan Y., Liu M., Guo X., Yang H. & Liu J. Guinea pig model for evaluating the potential public health risk of swine and avian influenza viruses. *PLoS One*, 2010, 5, e15537.
- Sürth V., Lopes de Carvalho I., Nuncio M. S., Cláudia Norte A. & Kraiczy P. Bactericidal activity of avian complement: a contribution to understand avian-host tropism of *Lyme borreliæ*. *Parasites and Vectors*, 2021, 14, 451.
- Taff C. C. & Townsend A. K. *Campylobacter jejuni* infection associated with relatively poor condition and low survival in a wild bird. *Journal of Avian Biology*, 2017, 48, 1071-6.
- Taheri N., Fällman M., Wai S. N. & Fahlgren A. Accumulation of virulence-associated proteins in *Campylobacter jejuni* outer membrane vesicles at human body temperature. *Journal of Proteomics*, 2019, 195, 33-40.
- Tsiouris V., Kiskinis K., Mantzios T., Dovas C. I., Mavromati N., Filiouis G., Brellou G., Vlemmas I. & Georgopoulou I. Avian mycobacteriosis and molecular identification of *Mycobacterium avium* Subsp. *avium* in racing pigeons (*Columba livia domestica*) in Greece. *Animals*, 2021, 11, 291.
- Ul-Rahman A., Ishaq H. M., Raza M. A. & Shabbir M. Z. Zoonotic potential of Newcastle disease virus: Old and novel perspectives related to public health. *Reviews in Medical Virology*, 2022, 32 (1), e2246.
- Venkatesan G., Balamurugan V., Gandhale P. N., Singh R. K. & Bhanuprakash V. Viral zoonosis: a comprehensive review. *Asian Journal of Animal and Veterinary Advances*, 2010, 5, 77-92.
- Wang Y., Niu S., Zhang B., Yang C. & Zhou Z. The whole genome analysis for the first human infection with H10N3 influenza virus in China. *Journal of Infection*, 2021, S0163-4453(21)00318-2.
- Wibisono F. M., Wibisono F. J., Effendi M. H., Plumeriastuti H., Hidayatullah A. R., Hartadi E. B. & Sofiana E. D. A Review of Salmonellosis on Poultry Farms: Public Health Importance. *Systematic Reviews in Pharmacy*, 2020, 11(9), 481-6.
- Wojtyniak B., Goryński P. & Moskalewicz B. Polish population health situation and its determinants. Report of PZH. *Warsaw*, 2013.
- Wright P., Neumann G. and Kawaoka Y. 2013, Orthomyxoviruses. In: *Fields virology*, Ed. D. Knipe and P. Howley. 6th ed. Lippincott Williams & Wilkins, Philadelphia, PA.
- Wu H. M., Huang W. Y., Lee M. L., Yang A. D., Chaou K. & Hsieh L. Clinical features, acute

- 
- complications, and outcome of Salmonella meningitis in children under one year of age in Taiwan. *BMC Infectious Diseases*, 2011, 11, 30.
- Yune N. & Abdela N. Update on Epidemiology, Diagnosis and Control Technique of Newcastle Disease. *Journal of Veterinary Science & Technology*, 2017, 8, 429.
- Zanaty A. M., Erfan A. M., Mady W. H., Amer F., Nour A. A., Rabie N., Samy M., Selim A. A., Hassan W. M. M. & Naguib M. M. Avian influenza virus surveillance in migratory birds in Egypt revealed a novel reassortant H6N2 subtype. *Avian Research*, 2019, 10, 1-10
- Zhang J., Li X., Wang X., Ye H., Li B., Chen Y., Chen J., Zhang T., Qiu Z., Li H., Jia W., Liao M. & Qi W. Genomic evolution, transmission dynamics, and pathogenicity of avian influenza A (H5N8) viruses emerging in China, 2020. *Virus Evolution*, 2021, 7(1), veab046.
- Zhang X. X., Zhang N. Z., Tian W. P., Zhou D. H., Xu Y. T., & Zhu X. Q. First report of *Toxoplasma gondii* seroprevalence in pet parrots in China. *Vector-Borne and Zoonotic Diseases*, 2014, 14(6), 394-8.
- Zhang Y., Cooper B., Gui X., Sherer R. & Cao Q. Clinical diversity of invasive cryptococcosis in AIDS patients from central China: report of two cases with review of literature. *BMC Infectious Diseases*, 2019, 19, 1003.
- Ziegler U., Lühken R., Keller M., Cadar D., Van der Grinten E., Michel F., Albrecht K., Eiden M., Rinder M., Lachmann L., Höper D., Vina-Rodriguez A., Gaede W., Pohl A., Schmidt-Chanasit J. & Groschup M. H. West Nile virus epizootic in Germany, 2018. *Antiviral Research*, 2019, 162, 39-43.
-