



Existence of solutions of Caputo fractional integro-differential equations

Manochehr Kazemi*

Department of Mathematics, Ashtian Branch, Islamic Azad University, Ashtian, Iran.

Abstract

In this paper, by using the techniques of measures of non-compactness and the Petryshyn fixed point theorem, we investigate the existence of solutions of a Caputo fractional functional integro-differential equation and obtain some new results. These existing results involve particular results gained from earlier studies under weaker conditions.

Keywords. Measures of non-compactness, Fractional functional integro-differential equation, Fixed point theorem, Existence of solutions.

2010 Mathematics Subject Classification. 47H09, 47H10.

1. INTRODUCTION

Kuratowski [22] introduced the concept of a measure of noncompactness (MNC). Recently, there have been several successful attempt to apply the concept of MNC in the study of the existence of solutions of nonlinear integral equations and integro-differential equations [5–8, 11, 12, 17, 18, 23–25, 31, 34, 35, 37]. The study integro-differential equations is linked to the wide applications of calculus in mathematical sciences. Therefore, they have received much attention. In recent years, many researchers have focused on the development of techniques for discussing the solutions of fractional differential equations and fractional integro-differential equations [3, 4, 9, 10, 14, 19, 26, 36].

Susahab *et al.* [33] investigated the existence of solutions for fractional integro-differential equations of the type

$$\begin{cases} {}^C D^\gamma(\chi(\vartheta)) = q(\vartheta) + \int_0^\vartheta k(\vartheta, \ell, \chi(\ell)) d\ell, & \vartheta \in [0, a], \\ \chi^{(i)}(0) = \chi_i, & i = 0, 1, \dots, n-1. \end{cases} \quad (1.1)$$

Karhikeyan *et al.* [16], studied the existence existence of solutions for fractional integro-differential equations of the type

$${}^C D^\gamma(\chi(\tau)) = f(\tau, \chi(\tau)) + \int_0^\tau k(\tau, s, \chi(s)) ds, \quad \tau \in [0, a]. \quad (1.2)$$

Recently, in [9], Dadsetadi and *et al.* established the existence and uniqueness of the solution of the following nonlinear fractional Volterra integro-differential equations with the help of Darbo's fixed point theorem,

$$\begin{cases} {}^C D^\gamma(\chi(\vartheta) + \rho(\vartheta, \chi(\vartheta))) = h(\vartheta, \chi(\vartheta)) + q\left(\vartheta, \chi(\vartheta), \int_0^\vartheta p(\vartheta, \ell) H(\chi(\ell)) d\ell\right), & \vartheta \in [0, a], \\ \chi^{(i)}(0) = \chi_i, & i = 0, 1, \dots, n-1. \end{cases} \quad (1.3)$$

In 2021, Samei *et al.* studied the following singular fractional integro-differential equation involving Caputo fractional q -derivative, for $0 < s < 1$,

$${}^C D_q^\sigma \chi(s) = g\left(s, \chi(s), \chi'(s), {}^C D_q^\zeta \chi(s), \int_0^s \chi(r) \beta(r) dr\right), \quad (1.4)$$

Received: 05 July 2023 ; Accepted: 03 August 2024.

* Corresponding author. Email: univer_ka@yahoo.com.

with conditions $\chi(0) = 0$ and $\chi(1) = {}^C D_{\zeta}^{\eta} \chi(\tau)$, where $\chi \in C([0, 1])$, $\sigma \in [1, 2)$, $\zeta, \eta, \tau \in (0, 1)$ [30]. Relevant results have been presented in other studies [1, 2, 15, 20, 28].

In the research, motivated by the above-mentioned achievements, we discuss new existing results for Caputo fractional Volterra integro-differential equation (FVIDE) of the form

$${}^C D^{\sigma}(\chi(\vartheta) + g(\vartheta, \chi(\vartheta))) = f(\vartheta, \chi(\alpha(\vartheta))) + F\left(\vartheta, \chi(\beta(\vartheta)), \frac{1}{\Gamma(\zeta)} \int_0^{\vartheta} \frac{k(\vartheta, \ell, \chi(\mu(\ell)))}{(\vartheta - \ell)^{1-\zeta}} d\ell\right), \quad \vartheta \in J_a = [0, a], \tag{1.5}$$

with the initial conditions

$$\chi^{(i)}(0) = \chi_i, \quad i = 0, 1, \dots, n - 1, \tag{1.6}$$

presented and proved. Here ${}^C D^{\sigma}$ is the Caputo’s fractional derivative and $\chi : J_a \rightarrow \mathcal{R}$ is an unknown function and other functions are known. Also $g : J_a \times \mathcal{R} \rightarrow \mathcal{R}$, $f : J_a \times \mathcal{R} \rightarrow \mathcal{R}$, $k : J_a^2 \times \mathcal{R} \rightarrow \mathcal{R}$, $\alpha, \beta, \mu : J_a \rightarrow J_a$ are continuous functions and $\chi_i, i = 0, 1, \dots, n - 1$, are constants. The main goal of this article is to investigate the new results on the existence of the solution of Eq. (1.5). For this, we use a fixed point theorem due to Petryshyn that has been presented as a generalization of Darbo’s fixed point theorem.

The paper is arranged as follows: In section 2, we recall some auxiliary facts and notations of the idea of MNC. Section 3 is involved in proving the main theorem for Eq. (1.5). In section 4, some examples are given to demonstrate the applicability of our results. Finally, section 5 concludes our work.

2. PRELIMINARIES

Definition 2.1 ([21]). The Riemann-Liouville fractional integral of order $\sigma > 0$ of a function χ , is defined as

$$I^{\sigma} \chi(\vartheta) = \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta} (\vartheta - \mu)^{\sigma-1} \chi(\mu) d\mu, \quad \vartheta > 0. \tag{2.1}$$

Definition 2.2 ([21]). The Caputo derivative of fractional order non negative $\sigma \geq 0$ for a function χ , is defined by

$$({}^C D^{\sigma} \chi)(\vartheta) = \frac{1}{\Gamma(n - \sigma)} \int_0^{\vartheta} (\vartheta - \mu)^{n-\sigma-1} \chi^{(n)}(\mu) d\mu, \quad n = [\sigma] + 1. \tag{2.2}$$

Lemma 2.3 ([21]). Let $\sigma > 0$ and $n = [\sigma] + 1$. If $\chi(\vartheta) \in C^n[0, a]$, then

$$(i) (I^{\sigma} {}^C D^{\sigma} \chi)(\tau) = \chi(\tau) - \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0)}{i!} \tau^i,$$

$$(ii) ({}^C D^{\sigma} I^{\sigma} \chi)(\tau) = \chi(\tau).$$

Throughout this paper, assume that $E = C([0, a])$ is a Banach space with the standard norm $\|\cdot\|$. Denote by $\bar{B}_{\delta} = \{z \in E : \|z\| \leq \delta\}$ the closed ball centered at the origin 0 of radius δ . The symbol $\partial \bar{B}_{\delta} = \{z \in E : \|z\| = \delta\}$ represents a sphere in E and around 0 with radius δ .

Definition 2.4 ([22]). Let $P \subset E$, so $\alpha(P)$ which is called Kuratowski MNC is

$$\alpha(P) = \inf \left\{ \sigma > 0 : P = \bigcup_{i=1}^n P_i, \text{diam}(P_i) \leq \sigma, i = 1, 2, \dots, n \right\}. \tag{2.3}$$

Definition 2.5 ([13]). Let $P \subset E$ and

$$\Delta(P) = \inf \left\{ \sigma > 0 : P \text{ has a finite } \sigma\text{-net in } E \right\}. \tag{2.4}$$

This quantity is called the Hausdorff MNC.

Theorem 2.6 ([29]). Let $P, Q \subset E$, then

- (i) $\Delta(P \cup Q) = \max\{\Delta(P), \Delta(Q)\}$,
- (ii) $\Delta(P + Q) \leq \Delta(P) + \Delta(Q)$,



- (iii) $\Delta(tP) = |t| \Delta(P)$, $t \in \mathcal{R}$,
- (iv) $\Delta(P) \leq \Delta(Q)$ for $P \subseteq Q$,
- (v) $\Delta(\bar{co}P) = \Delta(P)$,
- (vi) $\Delta(P)$ is zero if and only if P be precompact.

Theorem 2.7 ([13]). For all bounded sets P from space $C(J_a)$

$$\Delta(P) = \limsup_{\sigma \rightarrow 0} \sup_{y \in P} \omega^{J_a}(y, \sigma), \tag{2.5}$$

where $\omega^{J_a}(y, \sigma)$ is the modulus of continuity of function y on J_a .

Definition 2.8 ([27]). Let $\mathcal{U} : E \rightarrow E$ be a continuous mapping so that $\forall P \subset E$ with P bounded, $\mathcal{U}(P)$ is bounded and $\Delta(\mathcal{U}P) \leq \lambda \Delta(P)$, $\lambda \in (0, 1)$. If

$$\Delta(\mathcal{U}P) < \Delta(P), \quad \forall \Delta(P) > 0, \tag{2.6}$$

then \mathcal{U} is called condensing mapping.

Theorem 2.9 ([29], see also [32]). Let $\mathcal{U} : B_\delta \rightarrow E$ be a condensing mapping so that:

- (P) if $\mathcal{U}(x) = \lambda x$, for some x in ∂B_δ then $\lambda \leq 1$.

Then \mathcal{U} has at least one fixed point in B_δ .

3. MAIN RESULTS

In this section, fixed point theorems are used to check the existence of the solution to FVIDE (1.5). Applying Lemma 2.3, we get

$$\begin{aligned} \chi(\vartheta) &= \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i - g(\vartheta, \chi(\vartheta)) \\ &+ \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell, \end{aligned} \tag{3.1}$$

where

$$(H\chi)(\vartheta) = \frac{1}{\Gamma(\varsigma)} \int_0^\vartheta \frac{k(\vartheta, \ell, \chi(\mu(\ell)))}{(\vartheta - \ell)^{1-\varsigma}} d\ell. \tag{3.2}$$

The Eq. (1.5) is equivalent to the above fractional integral equation. So, every solution of (3.1) is a solution of (1.5) and vice versa. In what follows, we consider Eq. (3.1) under the following conditions:

- (L1) $g \in C(J_a \times \mathcal{R}, \mathcal{R})$, $f \in C(J_a \times \mathcal{R}, \mathcal{R})$, $F \in C(J_a \times \mathcal{R}^2, \mathcal{R})$, $k \in C(J_a^2 \times \mathcal{R}, \mathcal{R})$, and $\alpha, \beta, \mu : J_a \rightarrow J_a$ are continuous;
- (L2) There exist non negative constants k_1, k_2, c_1, c_2 , and c_3 so that $k_1 < 1$ and

$$\begin{aligned} |g(\vartheta, \omega_1) - g(\vartheta, \varpi_1)| &\leq k_1 |\omega_1 - \varpi_1|, \\ |F(\vartheta, \omega_1, \omega_2) - F(\vartheta, \varpi_1, \varpi_2)| &\leq c_1 |\omega_1 - \varpi_1| + c_2 |\omega_2 - \varpi_2|; \end{aligned} \tag{3.3}$$

- (L3) $\exists \delta_0 \geq 0$ such that

$$\sup \left\{ L + A + \frac{M_1 a^\sigma}{\Gamma(1 + \sigma)} + \frac{M_2 a^\sigma}{\Gamma(1 + \sigma)} \right\} \leq \delta_0, \tag{3.4}$$



where,

$$\begin{aligned}
 L &= \sup \left\{ \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i \right| : \forall \vartheta \in J_a \right\}, \\
 A &= \sup \left\{ |g(\vartheta, \omega_1)| : \forall \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}, \\
 M_1 &= \sup \left\{ |f(\vartheta, \omega_1)| : \forall \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}, \\
 M_2 &= \sup \left\{ |F(\vartheta, \omega_1, \omega_2)| : \forall \vartheta \in J_a, \omega_1, |\omega_2| \leq \frac{\vartheta^\varsigma B}{\Gamma(1 + \varsigma)} \right\}, \\
 B &= \sup \left\{ |k(\vartheta, \ell, \omega_1)| : \forall \vartheta, \ell \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}.
 \end{aligned} \tag{3.5}$$

Theorem 3.1. *With the conditions (L1)-(L3), Eq. (1.5) with the initial conditions (1.6), has at least one solution in $E = C(J_a)$.*

Proof. First, $\mathcal{U} : B_{\delta_0} \rightarrow E$ is define as follows

$$\begin{aligned}
 (\mathcal{U}\chi)(\vartheta) &= \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i - g(\vartheta, \chi(\vartheta)) \\
 &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell,
 \end{aligned} \tag{3.6}$$

then it is checked that \mathcal{U} is continuous on B_{δ_0} . Considering $\varepsilon > 0$ and for arbitrary values $\chi, \eta \in B_{\delta_0}$ such that $\|\chi - \eta\| \leq \varepsilon$, when $\vartheta \in J_a$ we will have

$$\begin{aligned}
 |(\mathcal{U}\chi)(\vartheta) - (\mathcal{U}\eta)(\vartheta)| &\leq |g(\vartheta, \chi(\vartheta)) - g(\vartheta, \eta(\vartheta))| \\
 &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{|f(\ell, \chi(\alpha(\ell))) - f(\ell, \eta(\alpha(\ell)))|}{(\vartheta - \ell)^{1-\sigma}} d\ell \\
 &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{|F(\ell, \chi(\beta(\ell)), (H\chi)(\ell)) - F(\ell, \eta(\beta(\ell)), (H\eta)(\ell))|}{(\vartheta - \ell)^{1-\sigma}} d\ell \\
 &\leq k_1 \|\chi - \eta\| + \frac{\vartheta^\sigma}{\Gamma(1 + \sigma)} \omega(f, \omega(\alpha, \varepsilon)) \\
 &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{|F(\ell, \chi(\beta(\ell)), (H\chi)(\ell)) - F(\ell, \eta(\beta(\ell)), (H\chi)(\ell))|}{(\vartheta - \ell)^{1-\sigma}} d\ell \\
 &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{|F(\ell, \eta(\beta(\ell)), (H\chi)(\ell)) - F(\ell, \eta(\beta(\ell)), (H\eta)(\ell))|}{(\vartheta - \ell)^{1-\sigma}} d\ell \\
 &\leq k_1 \|\chi - \eta\| + \frac{\vartheta^\sigma}{\Gamma(1 + \sigma)} \omega(f, \omega(\alpha, \varepsilon)) + \frac{c_1 \vartheta^\sigma}{\Gamma(1 + \sigma)} \|\chi - \eta\| + \frac{c_2 \vartheta^\sigma s^\varsigma}{\Gamma(1 + \sigma)\Gamma(1 + \varsigma)} \omega(k, \varepsilon),
 \end{aligned}$$

where

$$\begin{aligned}
 \omega(f, \omega(\alpha, \varepsilon)) &= \sup \left\{ |f(\ell, \chi) - f(\ell, \eta)| : \ell \in J_a, \chi, \eta \in [-\delta_0, \delta_0], \|\chi - \eta\| \leq \omega(\alpha, \varepsilon) \right\}, \\
 \omega(\alpha, \varepsilon) &= \sup \left\{ |\alpha(\ell_1) - \alpha(\ell_2)| : \ell_1, \ell_2 \in J_a, \|\ell_1 - \ell_2\| \leq \varepsilon \right\}, \\
 \omega(k, \varepsilon) &= \sup \left\{ |k(\vartheta, \ell, \chi) - k(\vartheta, \ell, \eta)| : \vartheta, \ell \in J_a, \chi, \eta \in [-\delta_0, \delta_0], \|\chi - \eta\| \leq \varepsilon \right\}.
 \end{aligned} \tag{3.7}$$

Now, because the functions $f = f(\ell, \chi)$ and $k = k(\vartheta, \ell, \chi)$ are uniformly continuous on $J_a \times \mathcal{R}$ and $J_a^2 \times \mathcal{R}$, respectively, we conclude $\omega(f, \omega(\alpha, \varepsilon)) \rightarrow 0$ and $\omega(k, \varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0$. Hence, the continuity of \mathcal{U} on B_{δ_0} results. Now, it is shown



\mathcal{U} satisfies densifying condition. Let ε be an arbitrary positive constant. For $\chi \in P \subset E$ let $\vartheta_1, \vartheta_2 \in J_a$ while $\vartheta_1 \leq \vartheta_2$ and $\vartheta_2 - \vartheta_1 \leq \varepsilon$. Therefore we obtain

$$\begin{aligned} |(\mathcal{U}\chi)(\vartheta_2) - (\mathcal{U}\chi)(\vartheta_1)| &= \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta_2^i - g(\vartheta_2, \chi(\vartheta_2)) \right. \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_2} \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_2} \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell \\ &\quad - \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta_1^i + g(\vartheta_1, \chi(\vartheta_1)) \\ &\quad - \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_1} \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_1 - \ell)^{1-\sigma}} d\ell - \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_1} \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_1 - \ell)^{1-\sigma}} d\ell \left. \right| \\ &\leq \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} (\vartheta_2^i - \vartheta_1^i) \right| \\ &\quad + |g(\vartheta_1, \chi(\vartheta_1)) - g(\vartheta_1, \chi(\vartheta_2))| + |g(\vartheta_1, \chi(\vartheta_2)) - g(\vartheta_2, \chi(\vartheta_2))| \\ &\quad + \frac{1}{\Gamma(\sigma)} \left| \int_0^{\vartheta_1} \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell + \int_{\vartheta_1}^{\vartheta_2} \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell - \int_0^{\vartheta_1} \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_1 - \ell)^{1-\sigma}} d\ell \right| \\ &\quad + \frac{1}{\Gamma(\sigma)} \left| \int_0^{\vartheta_1} \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell + \int_{\vartheta_1}^{\vartheta_2} \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_2 - \ell)^{1-\sigma}} d\ell \right. \\ &\quad \left. - \int_0^{\vartheta_1} \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(s))}{(\vartheta_1 - \ell)^{1-\sigma}} d\ell \right| \\ &\leq k_1 |\chi(\vartheta_1) - \chi(\vartheta_2)| + \omega_g(J_a, \varepsilon) \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_1} \left| \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_2 - \ell)^{1-\sigma}} - \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_1 - \ell)^{1-\sigma}} \right| d\ell + \frac{1}{\Gamma(\sigma)} \int_{\vartheta_1}^{\vartheta_2} \left| \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta_2 - \ell)^{1-\sigma}} \right| d\ell \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^{\vartheta_1} \left| \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_2 - \ell)^{1-\sigma}} - \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta_1 - \ell)^{1-\sigma}} \right| d\ell \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_{\vartheta_1}^{\vartheta_2} \left| \frac{F(\ell, \chi(\beta(\ell)), \chi(\varsigma(\ell)), (H\chi)(\ell))}{(\vartheta_2 - \ell)^{1-\sigma}} \right| d\ell. \end{aligned}$$

For simplicity, we use the following notation:

$$\omega_g(J_a, \varepsilon) = \sup \left\{ |g(\vartheta, \omega_1) - g(\bar{\vartheta}, \omega_1)| : |\vartheta - \bar{\vartheta}| \leq \varepsilon, \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}. \tag{3.8}$$

Then we have

$$\begin{aligned} |(\mathcal{U}\chi)(\vartheta_2) - (\mathcal{U}\chi)(\vartheta_1)| &\leq k_1 \omega(\chi, \varepsilon) + \omega_g(J_a, \varepsilon) \\ &\quad + \frac{M_1}{\Gamma(1 + \sigma)} \{ \vartheta_1^\sigma - \vartheta_2^\sigma + (\vartheta_2 - \vartheta_1)^\sigma \} + \frac{M_1}{\Gamma(1 + \sigma)} (\vartheta_2 - \vartheta_1)^\sigma \\ &\quad + \frac{M_2}{\Gamma(1 + \sigma)} \{ \vartheta_1^\sigma - \vartheta_2^\sigma + (\vartheta_2 - \vartheta_1)^\sigma \} + \frac{M_2}{\Gamma(1 + \sigma)} (\vartheta_2 - \vartheta_1)^\sigma \\ &\leq k_1 \omega(\chi, \varepsilon) + \omega_g(J_a, \varepsilon) + \frac{3\varepsilon^\sigma M_1}{\Gamma(1 + \sigma)} + \frac{3\varepsilon^\sigma M_2}{\Gamma(1 + \sigma)}. \end{aligned} \tag{3.9}$$

This yields the following estimate

$$\omega(\mathcal{U}\chi, \varepsilon) \leq k_1 \omega(\chi, \varepsilon), \quad \chi \in P. \tag{3.10}$$



Thus, taking the supremum in P , then the limit as $\varepsilon \rightarrow 0$, we obtain $\Delta(\mathcal{U}P) \leq k_1\Delta(P)$. Hence \mathcal{U} is a condensing map. It remains to verify condition (P) of Theorem 2.9. Let $\chi \in \partial\bar{B}_{\delta_0}$. If $\mathcal{U}\chi = \lambda\chi$ then we have $\lambda\delta_0 = \lambda\|\chi\| = \|\mathcal{U}\chi\|$, and with the condition (L3), we get

$$\begin{aligned} |\mathcal{U}\chi(\vartheta)| &= \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i - g(\vartheta, \chi(t)) \right. \\ &\quad \left. + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{F(\ell, \chi(\beta(\ell)), (H\chi)(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell \right| \leq \delta_0, \quad \vartheta \in J_a, \end{aligned} \tag{3.11}$$

and hence $\|\mathcal{U}\chi\| \leq \delta_0$, this means $\lambda \leq 1$. □

The following equation which is the main results of Dadsetadi and *et al.* [9], would be obtained from Theorem 3.1.

$${}^C D^\gamma(x(\vartheta) + g(\vartheta, x(\vartheta))) = f(\vartheta, x(\vartheta)) + F\left(\vartheta, x(\vartheta), \int_0^\vartheta q(\vartheta, \ell)H(x(\ell)) d\ell\right), \quad \vartheta \in J_a = [0, a], \tag{3.12}$$

with the initial conditions

$$x^{(i)}(0) = x_i, \quad i = 0, 1, \dots, n - 1, \tag{3.13}$$

Remark 3.2. By employing Riemann-Liouville fractional integrating and Lemma 2.3, Eq. (3.12) changes into

$$\begin{aligned} x(t) &= \sum_{i=0}^{n-1} \frac{x^{(i)}(0) + g^{(i)}(0, x_0)}{i!} \vartheta^i - g(\vartheta, x(\vartheta)) \\ &\quad + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{f(\ell, x(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell + \frac{1}{\Gamma(\sigma)} \int_0^\vartheta \frac{F(\ell, x(\ell), (\psi x)(\ell))}{(\vartheta - \ell)^{1-\sigma}} d\ell. \end{aligned} \tag{3.14}$$

where $(\psi x)(\vartheta) = \int_0^\vartheta q(\vartheta, \ell)H(x(\ell))d\ell$.

Corollary 3.3. Consider Eq. (3.14) under the following conditions

(M1) There exist non negative constants k_1, k_2, c_1, c_2 , and c_3 so that $k_1 < 1$ and

$$\begin{aligned} |g(\vartheta, \omega_1) - g(\vartheta, \varpi_1)| &\leq k_1|\omega_1 - \varpi_1| \\ |F(\vartheta, \omega_1, \omega_2) - F(\vartheta, \varpi_1, \varpi_2)| &\leq c_1|\omega_1 - \varpi_1| + c_2|\omega_2 - \varpi_2|, \end{aligned} \tag{3.15}$$

(M1) $\exists \delta_0 \geq 0$ such that

$$\sup \left\{ L + A + \frac{M_1 a^\sigma}{\Gamma(1 + \sigma)} + \frac{M_2 a^\sigma}{\Gamma(1 + \sigma)} \right\} \leq \delta_0, \tag{3.16}$$

where

$$\begin{aligned} L &= \sup \left\{ \left| \sum_{i=0}^{n-1} \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0)}{i!} \vartheta^i \right| : \forall \vartheta \in J_a \right\}, \\ A &= \sup \left\{ |g(\vartheta, \omega_1)| : \forall \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}, \\ M_1 &= \sup \left\{ |f(\vartheta, \omega_1)| : \forall \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0] \right\}, \\ M_2 &= \sup \left\{ |F(\vartheta, \omega_1, \omega_2)| : \forall \vartheta \in J_a, \omega_1 \in [-\delta_0, \delta_0], |\omega_2| \leq aB \right\}, \\ B &= \sup \left\{ |q(\vartheta, \ell)H(x(\ell))| : \forall \vartheta, \ell \in J_a, x \in [-\delta_0, \delta_0] \right\}. \end{aligned} \tag{3.17}$$

Then Eq. (3.12) has at least a solution in J_a .

Proof. It is clear that Eq. (3.12) is a particular case of Eq. (1.5). Here $\alpha(\vartheta) = \beta(\vartheta) = \mu(\vartheta) = \vartheta, \varsigma = 1$, and $k(\vartheta, \ell, x(\mu(\ell))) = q(\vartheta, \ell)H(x(\ell))$. The proof is done similarly to Theorem 3.1. □



4. EXAMPLES

Example 4.1. Consider

$${}^C D^{5/3} \left(\chi(\vartheta) + \frac{1}{3} e^{-\vartheta} \sin(\chi(\vartheta)) \right) = \frac{\vartheta^2}{3(1+\vartheta^2)} + \frac{(1+2\vartheta) \ln(1+|\chi(\sqrt{\vartheta})|)}{7} e^{-\vartheta} + \frac{\vartheta \sin(\chi(1-\vartheta))}{3} \\ + \frac{\sqrt{\vartheta}}{4+4\sqrt{\vartheta}} \frac{1}{\Gamma(1/2)} \int_0^\vartheta \frac{\vartheta e^{-\ell} \left(\frac{1}{3} + \int_0^\ell (\sin(\chi(\zeta^2)) + \frac{1}{2} \arctan(\frac{\chi(\zeta^2)}{1+\xi(\zeta^2)})) d\zeta \right)}{(\vartheta-\ell)^{1/2}} d\ell, \quad (4.1)$$

for $\vartheta \in J_a$, with

$$\chi^{(i)}(0) = \chi_i, \quad i = 0, 1. \quad (4.2)$$

Here, Eq. (4.1) is a particular case of Eq. (1.5) with $\sigma = \frac{5}{3}$, $n = 2$, $a = 1$, and

$$g(\vartheta, \omega_1) = \frac{1}{3} e^{-\vartheta} \sin(\omega_1), \\ f(\vartheta, \chi(\alpha(\vartheta))) = \frac{\vartheta^2}{3(1+\vartheta^2)} + \frac{(1+2\vartheta) \ln(1+|\chi(\sqrt{\vartheta})|)}{7} e^{-\vartheta}, \\ F(\vartheta, \omega_1, \omega_2) = \frac{\vartheta \sin(\omega_1)}{3} + \frac{\sqrt{\vartheta}}{4+4\sqrt{\vartheta}} \omega_2, \\ \omega_2 = \frac{1}{\Gamma(\frac{1}{2})} \int_0^\vartheta \frac{k(\vartheta, \ell, \chi(\mu(\ell)))}{(\vartheta-\ell)^{\frac{1}{2}}} d\ell, \\ k(\vartheta, \ell, \chi(\mu(\ell))) = \vartheta e^{-\ell} \left(\frac{1}{3} + \chi(\mu(\ell)) \right), \\ \chi(\mu(\ell)) = \int_0^\ell \left(\sin(\chi(\zeta^2)) + \frac{1}{2} \arctan \left(\frac{\chi(\zeta^2)}{1+\xi(\zeta^2)} \right) \right) d\zeta.$$

It is clear that (L1) holds. Also, conditions (L2) and (L3) are satisfied. we have

$$|g(\vartheta, \omega_1) - g(\vartheta, \varpi_1)| \leq \frac{1}{3} |\omega_1 - \varpi_1|, \quad (4.3)$$

and

$$|F(\vartheta, \omega_1, \omega_2) - F(\vartheta, \varpi_1, \varpi_2)| \leq \frac{1}{3} |\omega_1 - \varpi_1| + \frac{1}{4} |\omega_2 - \varpi_2|. \quad (4.4)$$

Here $k_1 = \frac{1}{3} < 1$, $c_1 = \frac{1}{3}$, $c_2 = \frac{1}{4}$. Also, suppose that $\|\chi\| \leq \delta_0$, $\delta_0 > 0$ and $\chi_0 = 0$, $\chi_1 = 1$, then we have

$$|\chi(\vartheta)| = \left| \sum_{i=0}^1 \frac{\chi^{(i)}(0) + g^{(i)}(0, \chi_0, 0)}{i!} \vartheta^i - \left(\frac{1}{3} e^{-\vartheta} \sin(\chi(\vartheta)) \right) \right. \\ \left. + \frac{1}{\Gamma(5/3)} \int_0^\vartheta \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta-\ell)^{2/3}} d\ell + \frac{1}{\Gamma(5/3)} \int_0^\vartheta \frac{F(\ell, \chi(\beta(\ell)), \chi(\zeta(\ell)), (H\chi)(\ell))}{(\vartheta-\ell)^{2/3}} d\ell \right| \\ \leq \frac{5}{3} + \frac{1}{\Gamma(5/3+1)} \left(\frac{1}{3} + \frac{3}{7} \delta_0 \right) + \frac{1}{\Gamma(5/3+1)} \left(\frac{1}{3} + \frac{1}{4\Gamma(1.5)} \left(\frac{4}{3} + \frac{\delta_0}{2} \right) \right), \quad \forall \vartheta \in J_a.$$

So, condition (L3) holds if

$$\frac{5}{3} + \frac{1}{\Gamma(5/3+1)} \left(\frac{1}{3} + \frac{3}{7} \delta_0 \right) + \frac{1}{\Gamma(5/3+1)} \left(\frac{1}{3} + \frac{1}{4\Gamma(1.5)} \left(\frac{4}{3} + \frac{\delta_0}{2} \right) \right) \leq \delta_0. \quad (4.5)$$

This shows that $\delta_0 = 3.7974$ is a solution of the above inequality. In view of Theorem 3.1, every problem (4.1) and (4.2) has at least one solution defined on $[0, 1]$.



Example 4.2. Consider

$$\begin{aligned}
 {}^c D^{0.5} \left(\chi(\vartheta) + \frac{1 + \sqrt{1 + \vartheta}}{4} + \frac{\vartheta \ln(1 + \chi(\vartheta))}{2} \right) &= \frac{1}{2} \vartheta e^{-\vartheta^2} + \frac{\sin(\vartheta) \chi(\vartheta \sqrt[3]{\vartheta})}{6(1 + \vartheta)} + \frac{|\chi(\sqrt{\vartheta})|}{6(1 + |\chi(\sqrt{\vartheta})|)} \\
 &+ \frac{1}{7(e^\vartheta + \vartheta)} e^{-\vartheta} \frac{1}{\Gamma(3/4)} \int_0^\vartheta \frac{(e^{-\vartheta} + \ell \vartheta \cos(\vartheta)) \left(\int_0^\ell \frac{\vartheta \sin(\chi(\vartheta))}{4} + \vartheta \left(1 + \frac{1}{9} \chi(\zeta^3) \right) d\zeta \right)}{(\vartheta - \ell)^{1/4}} d\ell,
 \end{aligned} \tag{4.6}$$

for $\vartheta \in [0, 1]$, with

$$\chi(0) = \chi_0 = 0. \tag{4.7}$$

In view of Eq. (1.5), we have $\sigma = 0.5$, $n = a = 1$, and

$$\begin{aligned}
 g(\vartheta, \chi(\vartheta)) &= \frac{1 + \sqrt{1 + \vartheta}}{4} + \frac{\vartheta \ln(1 + \chi(\vartheta))}{2}, \\
 f(\vartheta, \chi(\alpha(\vartheta))) &= \frac{1}{2} \vartheta e^{-\vartheta^2} + \frac{\sin(\vartheta) \chi(\vartheta \sqrt[3]{\vartheta})}{6(1 + \vartheta)}, \\
 F(t, \omega_1, \omega_2) &= \frac{|\omega_1|}{6(1 + |\omega_1|)} + \frac{1}{7(e^\vartheta + \vartheta)} e^{-\vartheta} \omega_2, \\
 \omega_2 &= \frac{1}{\Gamma(3/4)} \int_0^\vartheta \frac{k(\vartheta, \ell, \chi(\mu(\ell)))}{(\vartheta - \ell)^{1/4}} d\ell, \\
 k(\vartheta, \ell, \chi(\mu(\ell))) &= (e^{-\vartheta} + \ell \vartheta \cos(\vartheta)) \chi(\mu(\ell)), \\
 \chi(\mu(\ell)) &= \int_0^\ell \frac{\vartheta \sin(\chi(\vartheta))}{4} + \vartheta \left(1 + \frac{1}{9} \chi(\zeta^3) \right) d\zeta.
 \end{aligned}$$

Observe that (L1) holds. We show that conditions (L2) and (L3) are satisfied. So,

$$|g(\vartheta, \omega_1) - g(\vartheta, \varpi_1)| \leq \frac{1}{2} |\omega_1 - \varpi_1|, \tag{4.8}$$

and

$$|F(\vartheta, \omega_1, \omega_2) - F(\vartheta, \varpi_1, \varpi_2)| \leq \frac{1}{6} |\omega_1 - \varpi_1| + \frac{1}{7} |\omega_2 - \varpi_2|. \tag{4.9}$$

Here $k_1 = \frac{1}{2} < 1$, $c_1 = \frac{1}{6}$, $c_2 = \frac{1}{7}$. Also, suppose that $\|\chi\| \leq \delta_0$, $\delta_0 > 0$ and $\chi_0 = 0$, then we have

$$\begin{aligned}
 |\chi(\vartheta)| &= \left| \chi(0) + g(0, \chi_0) - g(\vartheta, \chi(\vartheta)) + \frac{1}{\Gamma(1/2)} \int_0^\vartheta \frac{f(\ell, \chi(\alpha(\ell)))}{(\vartheta - \ell)^{1/2}} d\ell \right. \\
 &\quad \left. + \frac{1}{\Gamma(1/2)} \int_0^\vartheta \frac{F(\ell, \chi(\beta(\ell)), \chi(\varsigma(\ell)), (H\chi)(\ell))}{(\vartheta - \ell)^{1/2}} d\ell \right| \\
 &\leq \left(\frac{5}{4} + \frac{1}{2} \delta_0 \right) + \frac{1}{\Gamma(1.5)} \left(\frac{1}{2} + \frac{1}{6} \delta_0 \right) + \frac{1}{\Gamma(1.5)} \left(\frac{1}{6} + \frac{2}{7\Gamma(1.75)} \left(\frac{5}{4} + \frac{1}{6} \delta_0 \right) \right) < \delta_0,
 \end{aligned} \tag{4.10}$$

for $\vartheta \in J_a$. This shows $\delta_0 = 8.2528$ is a solution of the above inequality. In view of Theorem 3.1, every problem (4.6)-(4.7) has at least one solution defined on $[0, 1]$.

5. CONCLUSION

This article handles the existence of solutions of a Caputo fractional functional integro-differential Equation (1.5) in the Banach space. With the assistance of the appropriate measure of noncompactness and the Petryshyn fixed point theorem hypothesis, we demonstrated our current results of the studied problem in Banach algebra. We presented some examples to illustrate the efficiency of our results.



REFERENCES

- [1] N. Adjimi, A. Boutiara, and M. E. Samei, *et al.*, *On solutions of a hybrid generalized Caputo-type problem via the noncompactness measure in the generalized version of Darbo's criterion*, Journal of Inequalities and Applications, 2023 (2023), 34.
- [2] P. Amiri and M. E. Samei, *Existence of Urysohn and Atangana-Baleanu fractional integral inclusion systems solutions via common fixed point of multi-valued operators*, Chaos, Solitons & Fractals, 165 (2022), 112822.
- [3] R. P. Agarwal, Y. Zhou, and Y. He, *Existence of fractional neutral functional differential equations*, Computers & Mathematics with Applications, 59(3) (2010), 1095–1100.
- [4] A. Anguraj, P. Karthikeyan, and J. J. Trujillo, *Existence of solutions to fractional mixed integrodifferential equations with nonlocal initial condition*, Advances in Difference Equations, 2011 (2011), 1–12.
- [5] R. Arab, *The existence of fixed points via the measure of noncompactness and its application to functional-integral equations*, Mediterr. J. Math., 13(2) (2016), 759–773.
- [6] R. Azimi, M. Mohagheghy Nezhad, and R. Pourgholi, *Legendre Spectral Tau Method for Solving the Fractional Integro-Differential Equations with A Weakly Singular Kernel*, Global Analysis and Discrete Mathematics, 4(4) (2022), 361–372.
- [7] C. Baishya, R. N. Premakumari, M. E. Samei, and M. K. Naik, *Chaos control of fractional order nonlinear Bloch equation by utilizing sliding mode controller*, Chaos, Solitons & Fractals, 174 (2023), 113773.
- [8] S. P. Bhairat and M. E. Samei, *Nonexistence of global solutions for a HilferKatugampola fractional differential problem*, Partial Differential Equations in Applied Mathematics, 7 (2023), 100495.
- [9] S. Dadsetadi, K. Nouri, and L. Torkzadeh, *Solvability of some nonlinear Integro-Differential equations of fractional order via measure of non-compactness*, The Pure and Applied Mathematics, 27(1) (2020), 13–24.
- [10] S. Dadsetadi and K. Nouri, *Study on existence of solution for some fractional integro differential equations via the iterative process*, Advances in Modelling and Analysis A, 55(2) (2018), 57–61.
- [11] A. Deep, Deepmala, and M. Rabbani, *A numerical method for solvability of some non-linear functional integral equations*, Appl. Math. Comput., 402 (2021), 125637.
- [12] A. M. El-Sayed, E. M. Hamdallah, and M. M. Ba-Ali, *Qualitative Study for a Delay Quadratic Functional Integro-Differential Equation of Arbitrary (Fractional) Orders*, Symmetry, 14(4) (2022), 748.
- [13] L. S. Goldenshtejn and A. S. Markus, *On the measure of non-compactness of bounded sets and of linear operators*, In Studies in Algebra and Math. Anal. (Russian), pages 45–54. Izdat. “Karta Moldovenjaske”, Kishinev, 1965.
- [14] F. Haddouchi, M. E. Samei, and S. Rezapour, *Study of a sequential ψ -Hilfer fractional integro-differential equations with nonlocal BCs*, J. Pseudo-Differ. Oper. Appl. 14 (2023), 61.
- [15] M. Houas and M. E. Samei, *Existence and Stability of Solutions for Linear and Nonlinear Damping of q -Fractional Duffing-Rayleigh Problem*, Mediterr. J. Math. 20 (2023), 148.
- [16] K. Karthikeyan and J. J. Trujillo, *Existence and Uniqueness Results for Fractional Integro-Differential Equations With Boundary Value Conditions*, Commun. Nonlinear Sci. Numer. Simulat, 17 (2012), 4037–4043.
- [17] M. Kazemi and R. Ezzati, *Existence of solution for some nonlinear two-dimensional volterra integral equations via measures of noncompactness*, Appl. Math. Comput., 275 (2016), 165–171.
- [18] M. Kazemi, *On existence of solutions for some functional integral equations in Banach algebra by fixed point theorem*, Int. J. Nonlinear Anal. Appl., 13 (2022), 451–466.
- [19] M. Kazemi, A. Deep, and A. R. Yaghoobnia, *Application of fixed point theorem on the study of the existence of solutions in some fractional stochastic functional integral equations*, Math. Sci., 18 (2024), 125–136.
- [20] T. Kherraz, M. Benbachir, M. Lakrib, M. E. Samei, M. K. Kaabar, and S. A. Bhanotar, *Existence and uniqueness results for fractional boundary value problems with multiple orders of fractional derivatives and integrals*, Chaos, Solitons & Fractals, 166 (2023), 113007.
- [21] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and applications of fractional differential equations*, Elsevier, 204 (2006).
- [22] K. Kuratowski, *Sur les espaces completes*, Fund. Math., 15 (1934), 301–335.
- [23] A. Lachouri, M. E. Samei, and A. Ardjouni, *Existence and stability analysis for a class of fractional pantograph q -difference equations with nonlocal boundary conditions*, Boundary Value Problems, 2023 (2023), 2.



- [24] L. M. Mishra and R. V. Agarwal, *On existence theorems for some nonlinear functional-integral equations*, Dynamic systems and Applications, *25*(3) (2016), 303–320.
- [25] H. K. Nashine and R. Arab, *Existence of solutions to nonlinear functional-integral equations via the measure of noncompactness*, J. Fixed Point Theory Appl., *20*(2) (2018), 66.
- [26] K. Nouri, M. Nazari, and B. Keramati, *Existence results for a coupled system of fractional integro-differential equations with time-dependent delay*, Journal of Fixed Point Theory and Applications, *19*(4) (2017), 2927–2943.
- [27] R. D. Nussbaum, *The fixed-point index and fixed point theorem for k -set contractions*, Pro Quest LLC, Ann Arbor, MI, 1969. Thesis (Ph.D.)—The University of Chicago.
- [28] P. R. Patle, M. Gabeleh, V. Rakočević, and M. E. Samei, *New best proximity point (pair) theorems via MNC and application to the existence of optimum solutions for a system of ψ -Hilfer fractional differential equations*, Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas, *117*(3) (2023), 124.
- [29] W. V. Petryshyn, *Structure of the fixed points sets of k -set-contractions*, Arch. Rational Mech. Anal., *40* (1970/1971), 312–328.
- [30] M. E. Samei, H. Zanganeh, and S. M. Aydogan, *Investigation of a class of the singular fractional integro-differential quantum equations with multi-step methods*, Journal of Mathematical Extension, *15*(15) (2021), 1–54.
- [31] M. E. Samei, *Employing Kuratowski measure of non-compactness for positive solutions of system of singular fractional q -differential equations with numerical effects*, Filomat, *34*(9) (2020), 2971–2989.
- [32] S. Singh, B. Watson, and P. Srivastava, *Fixed point theory and best approximation: the KKM-map principle*, Mathematics and its Applications, *424* (1997).
- [33] D. N. Susahab, S. Shahmorad, and M. Jahanshahi, *Efficient quadrature rules for solving nonlinear fractional integro-differential equations of the Hammerstein type*, Applied Mathematical Modelling, *39*(18) (2015), 5452–5458.
- [34] S. Unhaley and S. Kendre, *On existence and uniqueness results for iterative fractional integro-differential equation with deviating arguments*, Appl. Math. E-Notes, *19* (2019), 116–127.
- [35] C. Vetro and F. Vetro, *On the existence of at least a solution for functional integral equations via measure of noncompactness*, Banach J. Math. Anal., *11*(3) (2017), 497–512.
- [36] X. G. Yue, M. E. Samei, A. Fathipour, M. K. Kaabar, and A. Kashuri, *Using Krasnoselskii’s theorem to investigate the Cauchy and neutral fractional q -integro-differential equation via numerical technique*, Nonlinear Engineering, *11*(1) (2022), 186–206.
- [37] L. Zhang, B. Ahmad, G. Wang, and R. P. Agarwal, *Nonlinear fractional integro-differential equations on unbounded domains in a Banach space*, J. Comput. Appl. Math., *249* (2013), 51–56.

