



Investigating of highly dispersive solitons for concatenation model with power law nonlinearity using improved modified extended tanh-function method

Eman Salah¹, Islam Samir², Emad Abo El-Dahab³, Hamdy M. Ahmed¹, Medhat Ammar³, Mostafa Eslami⁴

¹Department of Physics and Engineering Mathematics, Higher Institute of Engineering, El-Shorouk Academy, El-Shorouk, Cairo, Egypt.

²Department of Mathematics and Engineering Physics, Faculty of Engineering, Aim Shams University, Cairo, Egypt.

³Department of Mathematics, Faculty of Science, Helwan University, Cairo, Egypt.

⁴Department of Mathematics, Faculty of Mathematical Sciences, University of Mazandaran, Babolsar, Iran.

Abstract

This research examines the phenomenon of optical solitons in the framework of the dispersive concatenation model, which incorporates three established models: the Lakshmanan-Porsezian-Daniel equation (LPDE), the Hirota equation (HE) and the nonlinear Schrödinger equation (NLSE). This model describe the soliton transmission dynamics across trans-continental and trans-oceanic dynamics. The model provided is situated within the context of nonlinear optics, a branch of optics that deals with optical phenomena in materials where the response of the medium to light is nonlinear. The equation appears to be a generalized model that combines several well-known equations from nonlinear optics. These equations often emerge as simplified descriptions of specific nonlinear effects in various optical systems. They capture phenomena like self-focusing, self-phase modulation, and soliton propagation, among others. The improved modified extended tanh scheme (IMETS) is utilized to derive solitons and other solutions for the investigated model. Many types of solutions are extracted with the help of the IMETS. These solutions including dark, bright, and singular solitons, Weierstrass elliptic and singular periodic solutions. The nature of the extracted solutions is illustrated by introducing both 2D and 3D graphical representations and setting the parameters with appropriate values.

Keywords. Bright solitons, Dark solitons, Exact solutions, Nonlinear partial differential equations.

2010 Mathematics Subject Classification. 35C07, 35C08, 35C09.

1. INTRODUCTION

A variety of physical systems such as fluid dynamics, optics and plasma physics are modeled with the aid of NLSE which characterizes the progression of a function with complex values that represent the wave envelope. The nonlinearity in the equation stems from the fact that the wave speed depends on the wave amplitude. The nonlinear Schrödinger equation is of utmost importance in examining optical solitons. As these solitons move through a nonlinear medium, they can maintain their velocities and shapes. NLSEs can be utilized for the purpose of designing and enhancing optical fiber systems for a wide range of applications, including telecommunications and optical computing.

Several recent studies have focused on soliton solutions including the Gerdjikov-Ivanov equation [1, 2], the Radhakrishnan-Kundu-Lakshmanan equation [3, 4], the Kundu-Eckhaus equation [5], HE [6], LPDE [7], and others. The dispersive concatenation model is a composite of three separate mathematical models: the NLSE with fifth order dispersion, the LPDE, and the HE model. Each of these models elucidates distinct physical phenomena associated with nonlinear waves, including optical solitons, rogue waves, and other nonlinear waves. By examining the concatenation model, scientists can gain a more comprehensive understanding of the characteristics of nonlinear waves and developing new solutions to the associated physical dilemmas. Researching the concatenation model is a potential area of study that

Received: 18 April 2024 ; Accepted: 12 November 2024.

* Corresponding author. Email: mostafa.eslami@umz.ac.ir.

can provide new insights into the mechanics of nonlinear waves [8–13]. The dispersive concatenation model reads as [14]:

$$\begin{aligned} i\varphi_t &+ a\varphi_{xx} + b|\varphi|^2\varphi - i\delta_1(\sigma_1\varphi_{xxx} + \sigma_2|\varphi|^2\varphi_x) + \delta_2[\sigma_3\varphi_{xxxx} + \sigma_4|\varphi|^2\varphi_{xx} \\ &+ \sigma_5|\varphi|^4\varphi + \sigma_6|\varphi_x|^2\varphi + \sigma_7(\varphi_x)^2\varphi^* + \sigma_8\varphi_{xx}^*\varphi^2] - i\delta_3[\sigma_9\varphi_{xxxxx} + \sigma_{10}|\varphi|^2\varphi_{xxx} \\ &+ \sigma_{11}|\varphi|^4\varphi_x + \sigma_{12}\varphi\varphi_x\varphi_{xx}^* + \sigma_{13}\varphi^*\varphi_x\varphi_{xx} + \sigma_{14}\varphi\varphi_x^*\varphi_{xx} + \sigma_{15}\varphi_x^*(\varphi_x)^2] = 0. \end{aligned} \quad (1.1)$$

Here $\varphi(x, t)$ describes the wave profile. The temporal evolution is denoted by the first term. Chromatic dispersion and Self-Phase modulation are denoted by a and b respectively. All parameters σ_j for $j = 1, 2, 3, \dots, 15$ are constants. Eq. (1.1) converts to the classical NLSE when $\delta_1 = \delta_2 = \delta_3 = 0$ while HE model is generated in case of $\delta_1 \neq 0$ and $\delta_2 = \delta_3 = 0$. For $\delta_1 = \delta_3 = 0$ with $\delta_2 \neq 0$, Eq. (1.1) reduces to LPDE.

The proposed model was studied by employing the enhanced Kudryashov's approach to get straddled, bright, and singular optical solitons. This algorithm not only provides a nuanced understanding of the various soliton types but also highlights the occurrence of singular solitons that exhibit unique characteristics. In addition, the Riccati equation expansion approach was applied to obtain dark solitons in addition to singular solitons. Furthermore, the Weierstrass' expansion scheme was implemented to encompass bright, singular solitons [14]. In this work, the improved modified extended tanh function method is implemented to study Eq. (1.1). This method is based on the extended Riccati equation which can produce various and novel types of solutions for the investigated model. This solutions including bright, dark and singular solitons. In addition, singular periodic, rational, exponential and Weierstrass elliptic solutions can be raised.

2. INTEGRATION SCHEME OF IMETS

This part briefly discusses the IMETF approach [15, 16]. Considering the next NLPDE

$$F(\varphi, \varphi_t, \varphi_x, \varphi_{xx}, \varphi_{tx}, \dots) = 0. \quad (2.1)$$

The subsequent steps should be performed to handle Eq. (2.1) using the IMETF scheme:

Step(1): Utilizing the subsequent transformation:

$$\varphi(x, t) = \varphi(\xi), \quad (2.2)$$

where $\xi = x - ct$. After that, Eq. (2.1) can be expressed as:

$$G(\varphi, \varphi', \varphi'', \varphi''', \dots) = 0. \quad (2.3)$$

Step(2): The resulted NLODE's solutions can be represented as:

$$\varphi(\xi) = \sum_{j=0}^k a_j \Upsilon^j(\xi) + \sum_{j=-1}^{-k} b_{-j} \Upsilon^j(\xi), \quad (2.4)$$

where $\Upsilon(\xi)$ satisfy the following differential equation:

$$\Upsilon'(\xi) = \sqrt{d_0 + d_1\Upsilon(\xi) + d_2\Upsilon^2(\xi) + d_3\Upsilon^3(\xi) + d_4\Upsilon^4(\xi)}. \quad (2.5)$$

Step(3): k can be evaluated by applying the balance rule on Eq. (2.3).

Step(4): The NLODE is transformed into nonlinear algebraic equation (NLAE) by inserting Eqs. (2.4) and (2.5) into Eq. (2.3).

Step(5): A system of NLAE is then generated by collecting and equating the coefficients of $\Upsilon^j(\xi)$ with zero. The resultant system can be handled to evaluate a_j , b_j , and c using Mathematica packages.

Step(6): One can get many solutions for Eq. (2.1) by substituting the determined constants and the general solutions of (2.5) into Eq. (2.4).

Comparing with other techniques such as extended direct algebraic method [17], simple ansatz method [18], the Unified method [19], the modified and extended rational expansion method [20], the extended (G'/G^2) -expansion technique [21], the variational principle method [22, 23], this method give various types of solutions such as bright,



dark, singular solitons. In addition, other mathematical solutions such as exponential, Weierstrass elliptic, rational type and singular periodic solutions can be extracted. However, when N , which is evaluated via the balance rule, is large, it will give a more complex system that is difficult to solve.

3. MATHEMATICAL ANALYSIS

The purpose of this section is to obtain exact solutions to Eq. (1.1) in the format indicated below:

$$\begin{aligned}\varphi(x, t) &= \Re(\xi)e^{i\psi(x,t)}, \\ \xi &= x - ct, \\ \psi(x, t) &= -\kappa x + \omega t + \theta_0.\end{aligned}\tag{3.1}$$

Here κ denote the soliton frequency while θ_0 denote the phase constant. Wave number is denoted by ω while c denote the soliton speed. By inserting Eq. (3.1) into Eq. (1.1) and then dividing into real (Re) and imaginary (IM) portions, we obtain:

Re parts:

$$\begin{aligned}(\delta_2\sigma_3 - 5\delta_3\kappa\sigma_9)\Re^{(4)}(\xi) + (10\delta_3\kappa^3\sigma_9 - 6\delta_2\kappa^2\sigma_3 - 3\delta_1\kappa\sigma_1 + a)\Re''(\xi) \\ + (\delta_1\kappa^3\sigma_1 + \delta_2\kappa^4\sigma_3 - \delta_3\kappa^5\sigma_9 - a\kappa^2 - \omega)\Re(\xi) \\ + (2\delta_3\kappa\sigma_{12} - 2\delta_3\kappa\sigma_{13} - 2\delta_3\kappa\sigma_{14} - \delta_3\kappa\sigma_{15} + \delta_2\sigma_6 + \delta_2\sigma_7)\Re(\xi)\Re'(\xi) \\ + (-3\delta_3\kappa\sigma_{10} - \delta_3\kappa\sigma_{12} - \delta_3\kappa\sigma_{13} + \delta_3\kappa\sigma_{14} + \delta_2\sigma_4 + \delta_2\sigma_8)\Re^2(\xi)\Re''(\xi) + (\delta_2\sigma_5 - \delta_3\kappa\sigma_{11})\Re^5(\xi) \\ + (\delta_3\kappa^3\sigma_{10} + \delta_3\kappa^3\sigma_{12} + \delta_3\kappa^3\sigma_{13} - \delta_3\kappa^3\sigma_{14} - \delta_3\kappa^3\sigma_{15} - \delta_1\kappa\sigma_2 - \delta_2\kappa^2\sigma_4 + \delta_2\kappa^2\sigma_6 - \delta_2\kappa^2\sigma_7 \\ - \delta_2\kappa^2\sigma_8 + b)\Re^3(\xi) = 0,\end{aligned}\tag{3.2}$$

and the IM parts:

$$\begin{aligned}-\delta_3\sigma_9\Re^{(5)}(\xi) + (10\delta_3\kappa^2\sigma_9 - 4\delta_2\kappa\sigma_3 - \delta_1\sigma_1)\Re'''(\xi) + (-5\delta_3\kappa^4\sigma_9 + 4\delta_2\kappa^3\sigma_3 + 3\delta_1\kappa^2\sigma_1 - 2a\kappa - c)\Re'(\xi) \\ - \delta_3\sigma_{15}\Re^3(\xi) - (\delta_3\sigma_{12} + \delta_3\sigma_{13} + \delta_3\sigma_{14})\Re(\xi)\Re'(\xi)\Re''(\xi) - \delta_3\sigma_{10}\Re^2(\xi)\Re'''(\xi) \\ + (3\delta_3\kappa^2\sigma_{10} - \delta_3\kappa^2\sigma_{12} + 3\delta_3\kappa^2\sigma_{13} - \delta_3\kappa^2\sigma_{14} - \delta_3\kappa^2\sigma_{15} - 2\delta_2\kappa\sigma_4 - 2\delta_2\kappa\sigma_7 \\ + 2\delta_2\kappa\sigma_8 - \delta_1\sigma_2)\Re^2(\xi)\Re'(\xi) - \delta_3\sigma_{11}\Re^4(\xi)\Re'(\xi) = 0.\end{aligned}\tag{3.3}$$

Differentiate Eq. (3.2) with regard to ξ , we get

$$\begin{aligned}(\delta_2\sigma_3 - 5\delta_3\kappa\sigma_9)\Re^{(5)}(\xi) + (10\delta_3\kappa^3\sigma_9 - 6\delta_2\kappa^2\sigma_3 - 3\delta_1\kappa\sigma_1 + a)\Re''''(\xi) \\ (\delta_1\kappa^3\sigma_1 + \delta_2\kappa^4\sigma_3 - \delta_3\kappa^5\sigma_9 - a\kappa^2 - \omega)\Re'(\xi) \\ (2\delta_3\kappa\sigma_{12} - 2\delta_3\kappa\sigma_{13} - 2\delta_3\kappa\sigma_{14} - \delta_3\kappa\sigma_{15} + \delta_2\sigma_6 + \delta_2\sigma_7)\Re^3(\xi) + (2\delta_3\kappa\sigma_{12} - 6\delta_3\kappa\sigma_{13} \\ - 2\delta_3\kappa\sigma_{14} - 2\delta_3\kappa\sigma_{15} + 2\delta_2\sigma_6 + 2\delta_2\sigma_7 - 6\delta_3\kappa\sigma_{10} + 2\delta_2\sigma_4 + 2\delta_2\sigma_8)\Re(\xi)\Re'(\xi)\Re''(\xi) \\ + (-3\delta_3\kappa\sigma_{10} - \delta_3\kappa\sigma_{12} - \delta_3\kappa\sigma_{13} + \delta_3\kappa\sigma_{14} + \delta_2\sigma_4 + \delta_2\sigma_8)\Re^2(\xi)\Re''''(\xi) + 5(\delta_2\sigma_5 - \delta_3\kappa\sigma_{11})\Re^4(\xi)\Re'(\xi) \\ + 3(\delta_3\kappa^3\sigma_{10} + \delta_3\kappa^3\sigma_{12} + \delta_3\kappa^3\sigma_{13} - \delta_3\kappa^3\sigma_{14} - \delta_3\kappa^3\sigma_{15} - \delta_1\kappa\sigma_2 - \delta_2\kappa^2\sigma_4 \\ + \delta_2\kappa^2\sigma_6 - \delta_2\kappa^2\sigma_7 - \delta_2\kappa^2\sigma_8 + b)\Re^2(\xi)\Re'(\xi) = 0.\end{aligned}\tag{3.4}$$



Eqs. (3.3) and (3.4) will be similar under the following circumstances:

$$\begin{aligned}
& \delta_2\sigma_3 + \delta_3\sigma_9(1 - 5\kappa) = 0, \\
& 10\delta_3\kappa^2\sigma_9(\kappa - 1) + a + 2\sigma_3\delta_2\kappa(2 - 3\kappa) + \delta_1\sigma_1(1 - 3\kappa) = 0, \\
& \delta_1\kappa^2\sigma_1(\kappa - 3) + \delta_2\sigma_3\kappa^3(\kappa - 4) - \delta_3\sigma_9\kappa^4(\kappa - 5) + a\kappa^2 - \omega + 2a\kappa + c = 0, \\
& 2\delta_3\kappa\sigma_{12} - 2\delta_3\kappa\sigma_{13} - 2\delta_3\kappa\sigma_{14} + \delta_3\sigma_{15}(1 - \kappa) + \delta_2\sigma_6 + \delta_2\sigma_7 = 0, \\
& \delta_3\sigma_{12}(2\kappa + 1) + \delta_3\sigma_{13}(1 - 6\kappa) + \delta_3\sigma_{14}(1 - 2\kappa) - 2\delta_3\kappa\sigma_{15} + 2\delta_2\sigma_6 + 2\delta_2\sigma_7 - 6\delta_3\kappa\sigma_{10} + 2\delta_2\sigma_4 + 2\delta_2\sigma_8 = 0, \\
& \delta_3\sigma_{10}(1 - 3\kappa) - \delta_3\kappa\sigma_{12} - \delta_3\kappa\sigma_{13} + \delta_3\kappa\sigma_{14} + \delta_2\sigma_4 + \delta_2\sigma_8 = 0, \\
& 5\delta_2\sigma_5 + \delta_3\sigma_{11}(1 - 5\kappa) = 0, \\
& 3\delta_3\kappa^2\sigma_{10}(\kappa - 1) + \delta_3\kappa^2\sigma_{12}(1 - 3\kappa) + 3\delta_3\sigma_{13}\kappa^2(\kappa - 1) + \delta_3\kappa^2\sigma_{14}(1 - 3\kappa)\delta_3\kappa^2\sigma_{15}(1 - 3\kappa) \\
& + \delta_1\sigma_2(1 - 3\kappa) + \delta_2\kappa\sigma_4(2 - 3\kappa) + 3\delta_2\kappa^2\sigma_6 + \delta_2\kappa\sigma_7(2 - 3\kappa) - \delta_2\kappa\sigma_8(3\kappa + 2) + 3b = 0.
\end{aligned}$$

The following form can now be used to rewrite Eq.(3.3):

$$\begin{aligned}
& \Re^{(5)}(\xi) + \Delta_1\Re'''(\xi) + \Delta_2\Re'(\xi) + \Delta_3\Re'^3(\xi) \\
& + \Delta_4\Re(\xi)\Re'(\xi)\Re''(\xi) + \Delta_5\Re^2(\xi)\Re'''(\xi) + \Delta_6\Re^2(\xi)\Re'(\xi) + \Delta_7\Re^4(\xi)\Re'(\xi) = 0,
\end{aligned} \tag{3.5}$$

where

$$\begin{aligned}
\Delta_1 &= \frac{-1}{\delta_3\sigma_9}(10\delta_3\kappa^2\sigma_9 - 4\delta_2\kappa\sigma_3 - \delta_1\sigma_1), \\
\Delta_2 &= \frac{-1}{\delta_3\sigma_9}(-5\delta_3\kappa^4\sigma_9 + 4\delta_2\kappa^3\sigma_3 + 3\delta_1\kappa^2\sigma_1 - 2a\kappa - c), \\
\Delta_3 &= \frac{\sigma_{15}}{\sigma_9}, \\
\Delta_4 &= \frac{1}{\sigma_9}(\sigma_{12} + \sigma_{13} + \sigma_{14}), \\
\Delta_5 &= \frac{\sigma_{10}}{\sigma_9}, \\
\Delta_6 &= \frac{-1}{\delta_3\sigma_9}(3\delta_3\kappa^2\sigma_{10} - \delta_3\kappa^2\sigma_{12} + 3\delta_3\kappa^2\sigma_{13} - \delta_3\kappa^2\sigma_{14} - \delta_3\kappa^2\sigma_{15} - 2\delta_2\kappa\sigma_4 - 2\delta_2\kappa\sigma_7 + 2\delta_2\kappa\sigma_8 - \delta_1\sigma_2), \\
\Delta_7 &= \frac{\sigma_{11}}{\sigma_9}.
\end{aligned}$$

Applying balance rule, we have $k = 1$. Subsequently, the solutions of Eq. (3.5) can be represented as:

$$\Re_1(\xi) = a_0 + a_1\Upsilon + \frac{b_1}{\Upsilon}. \tag{3.6}$$

Using the procedures (2.3), (2.4) and (2.5) mentioned in the last part, the next results are derived:

Set 1. $g_0 = g_1 = g_3 = 0$

Result (I)

$$\begin{aligned}
a_0 = 0, b_1 = 0, \Delta_2 &= -\frac{2g_4(a_1^2\Delta_6 + 6\Delta_1g_4)(a_1^4\Delta_1\Delta_7 + a_1^2(2\Delta_6 - \Delta_1(\Delta_3 - 4\Delta_5))g_4 + 12\Delta_1g_4^2)}{a_1^4((\Delta_3 - 4\Delta_5)g_4 - a_1^2\Delta_7)^2}, \\
g_2 &= \frac{2g_4(a_1^2\Delta_6 + 6\Delta_1g_4)}{a_1^2(a_1^2\Delta_7 - \Delta_3g_4 + 4\Delta_5g_4)}, g_4 = -\frac{2a_1^2\Delta_7}{\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7 + \Delta_3 + 2\Delta_4 + 6\Delta_5}}.
\end{aligned}$$



Then, we derive

$$\varphi(x, t) = \left\{ \sqrt{\frac{24\Delta_1\Delta_7 - 2\Delta_6 \left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5 \right)}{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + 3\Delta_3 + 2\Delta_4 - 2\Delta_5 \right) \Delta_7}} \right. \\ \left. \times \operatorname{sech} \left(\frac{2(ct - x) \sqrt{\frac{12\Delta_1\Delta_7}{\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5} - \Delta_6}}{\sqrt{\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + 3\Delta_3 + 2\Delta_4 - 2\Delta_5}} \right) \right\} \times e^{i\psi(x,t)}, \quad (3.7)$$

$$\varphi(x, t) = \left\{ \sqrt{\frac{24\Delta_1\Delta_7 - 2\Delta_6 \left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5 \right)}{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + 3\Delta_3 + 2\Delta_4 - 2\Delta_5 \right) \Delta_7}} \right. \\ \left. \times \operatorname{sec} \left(\frac{2(ct - x) \sqrt{\Delta_6 - \frac{12\Delta_1\Delta_7}{\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5}}}{\sqrt{\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + 3\Delta_3 + 2\Delta_4 - 2\Delta_5}} \right) \right\} \times e^{i\psi(x,t)}. \quad (3.8)$$

A bright solitary is denoted by Eq. (3.7) whereas Eq. (3.8) denote a singular periodic solution.

Set 2. $g_1 = g_3 = 0, g_0 = \frac{g_2^2}{4g_4}$

Result (I)

$$a_0 = 0, \quad b_1 = 0, \quad g_4 = -\frac{a_1^2 \Delta_3 g_2^2}{4(\Delta_2 + \Delta_1 g_2 + 4g_2^2)}, \\ \Delta_6 = \frac{a_1^2 \Delta_3 g_2^2 - 2a_1^2 \Delta_4 g_2^2 - 2a_1^2 \Delta_5 g_2^2 + 12\Delta_2 g_4 - 72g_4 g_2^2}{2a_1^2 g_2}, \\ \Delta_7 = -\frac{g_4 (a_1^2 \Delta_3 + 2a_1^2 \Delta_4 + 6a_1^2 \Delta_5 + 120g_4)}{a_1^4}.$$

Then, we derive

$$\varphi(x, t) = \sqrt{2} \sqrt{\frac{\Delta_2 + \Delta_1 g_2 + 4g_2^2}{\Delta_3 g_2}} \times \tanh \left(\frac{\sqrt{-g_2}(x - ct)}{\sqrt{2}} \right) \times e^{i\psi(x,t)}, \quad (3.9)$$

$$\varphi(x, t) = \sqrt{2} \sqrt{-\frac{\Delta_2 + \Delta_1 g_2 + 4g_2^2}{\Delta_3 g_2}} \times \tan \left(\frac{\sqrt{g_2}(x - ct)}{\sqrt{2}} \right) \times e^{i\psi(x,t)}. \quad (3.10)$$

A dark solitary is denoted by Eq. (3.9) whereas Eq. (3.10) denotes a singular periodic solution.

Result (II)

$$a_0 = 0, \quad g_4 = -\frac{a_1 g_2}{2b_1}, \quad b_1 = \frac{\Delta_2 + 4g_2 (\Delta_1 + 16g_2)}{8a_1 \Delta_3 g_2}, \\ a_1 = \frac{(\Delta_3 + 2\Delta_4 + 6\Delta_5) g_2 - \sqrt{((\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7) g_2^2}}{4\Delta_7}, \\ \Delta_6 = \frac{8g_2 (a_1 b_1 (\Delta_3 - 2(\Delta_4 + \Delta_5)) + 36g_2) - 3\Delta_2}{4a_1 b_1}.$$



Then, we derive

$$\varphi(x, t) = \frac{-1}{\sqrt{2}\sqrt{\Delta_3 g_2}} \left\{ \sqrt{\Delta_2 + 4g_2(\Delta_1 + 16g_2)} \times \coth \left(\sqrt{2}\sqrt{-g_2}(x - ct) \right) \right\} \times e^{i\psi(x, t)}, \quad (3.11)$$

$$\varphi(x, t) = \frac{1}{\sqrt{\Delta_3 g_2}} \left\{ \sqrt{-\frac{\Delta_2}{2} - 2\Delta_1 g_2 - 32g_2^2} \times \cot \left(\sqrt{2}\sqrt{g_2}(x - ct) \right) \right\} \times e^{i\psi(x, t)}. \quad (3.12)$$

Eq. (3.11) is a singular solitary solution whereas Eq. (3.12) represents a singular periodic solution.

Set (3). $g_2 = g_4 = 0$

Result (I)

$$\begin{aligned} b_1 &= 0, \quad \Delta_7 = 0, \quad \Delta_3 = \frac{1}{2}(-3)(\Delta_4 + 2\Delta_5), \\ \Delta_1 &= \frac{a_1^3 \Delta_4 g_1 + 3a_1^3 \Delta_5 g_1 + 3a_0^2 a_1 \Delta_4 g_3 + 9a_0^2 a_1 \Delta_5 g_3 + 45a_0 g_3^2}{3a_1 g_3}, \\ \Delta_2 &= \frac{3a_1^4 (\Delta_4 + 2\Delta_5) g_0 - a_0 a_1^3 \Delta_4 g_1 + 3a_0^3 a_1 (\Delta_4 + 4\Delta_5) g_3 - 9a_1^2 g_1 g_3 + 45a_0^2 g_3^2}{2a_1^2}, \\ g_3 &= -\frac{1}{30} a_1 \left(\sqrt{a_0^2 (\Delta_4 + 4\Delta_5)^2 - 40\Delta_6 + a_0 (\Delta_4 + 4\Delta_5)} \right). \end{aligned}$$

Then, we derive

$$\varphi(x, t) = \left\{ a_0 + a_1 \wp \left(\frac{(x - ct) \sqrt{-a_1 \left(\sqrt{a_0^2 (\Delta_4 + 4\Delta_5)^2 - 40\Delta_6 + a_0 (\Delta_4 + 4\Delta_5)} \right)}}{2\sqrt{30}}, \{h_2, h_3\} \right) \right\} \times e^{i\psi(x, t)}, \quad (3.13)$$

where $h_2 = -4g_1/g_3$ and $h_3 = -4g_0/g_3$. Eq. (3.13) denotes Weierstrass elliptic solution

Set (4). $g_0 = g_1 = g_2 = 0$

Result (I)

$$\begin{aligned} b_1 &= 0, \quad a_0 = -\frac{\sqrt{-\frac{\sqrt{\Delta_6^2 - 4\Delta_2 \Delta_7 + \Delta_6}}{\Delta_7}}}{\sqrt{2}}, \quad g_3 = \frac{4a_0 g_4}{a_1}, \quad g_4 = \frac{a_1^2 \Delta_7}{3(\Delta_3 - 2\Delta_5)}, \\ \Delta_1 &= \frac{-2a_1^2 a_0^2 \Delta_7 - a_1^2 \Delta_6 - 6a_0^2 \Delta_5 g_4}{6g_4}, \quad \Delta_4 = -\frac{2(a_1^4 \Delta_7 + 6a_1^2 \Delta_5 g_4 + 90g_4^2)}{3a_1^2 g_4}. \end{aligned}$$

Then, we derive

$$\varphi(x, t) = \left\{ \frac{\sqrt{-\frac{\sqrt{\Delta_6^2 - 4\Delta_2 \Delta_7 + \Delta_6}}{\Delta_7}} \left(-2(x - ct)^2 \left(\sqrt{\Delta_6^2 - 4\Delta_2 \Delta_7 + \Delta_6} \right) + 9\Delta_3 - 18\Delta_5 \right)}{\sqrt{2} \left(2(x - ct)^2 \left(\sqrt{\Delta_6^2 - 4\Delta_2 \Delta_7 + \Delta_6} \right) + 3\Delta_3 - 6\Delta_5 \right)} \right\} \times e^{i\psi(x, t)}, \quad (3.14)$$



$$\varphi(x, t) = \left\{ \sqrt{-\frac{\sqrt{\Delta_6^2 - 4\Delta_2\Delta_7} + \Delta_6}{2\Delta_7}} \times \left(-1 - 2e^{\sqrt{\frac{2}{3}}(x-ct)} \sqrt{\frac{\sqrt{\Delta_6^2 - 4\Delta_2\Delta_7} + \Delta_6}{\Delta_3 - 2\Delta_5}} \right) \right\} \times e^{i\psi(x,t)}. \quad (3.15)$$

A rational solution is derived by Eq. (3.14) whereas Eq. (3.15) denotes an exponential solution.

Set (5). $g_0 = g_1 = 0, g_4 = \frac{g_3^2}{4g_2}$

Result (I)

$$\begin{aligned} b_1 = 0, a_0 &= \frac{a_1 g_2}{g_3}, g_2 = -\frac{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5\right) g_3^2}{8a_1^2 \Delta_7}, \\ \Delta_1 &= -\frac{g_2 (4a_1^4 \Delta_7 g_2^2 + a_1^2 \Delta_4 g_2 g_3^2 + 5a_1^2 \Delta_5 g_2 g_3^2 + 2a_1^2 \Delta_6 g_3^2 + 15g_3^4)}{3g_3^4}, \\ \Delta_2 &= \frac{g_2^2 (a_1^4 \Delta_7 g_2^2 + a_1^2 \Delta_4 g_2 g_3^2 + 2a_1^2 \Delta_5 g_2 g_3^2 - a_1^2 \Delta_6 g_3^2 + 12g_3^4)}{3g_3^4}. \end{aligned}$$

Then, we derive dark solitary solution

$$\begin{aligned} \varphi(x, t) &= \left\{ \frac{1}{4\sqrt{2}} \times \sqrt{-\frac{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5\right) g_3^2}{\Delta_7 g_4}} \right. \\ &\times \left(\tanh \left(\frac{(x - ct) \sqrt{-\frac{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5\right) g_3^2}{a_1^2 \Delta_7}}}{4\sqrt{2}} \right) + 1 \right) \\ &\left. - \frac{\left(\sqrt{(\Delta_3 + 2\Delta_4 + 6\Delta_5)^2 - 480\Delta_7} + \Delta_3 + 2\Delta_4 + 6\Delta_5\right) g_3}{8a_1 \Delta_7} \right\} \times e^{i\psi(x,t)}. \quad (3.16) \end{aligned}$$

4. 3D AND 2D VISUALIZATIONS

In order to highlight the properties of the derived solutions, we present the two-dimensional and three-dimensional graphs of some solutions. Fig (1) shows a bright solitary solution of Eq. (3.7) with $c = 0, \Delta_1 = 2, \Delta_2 = 0.18, \Delta_3 = 0.25, \Delta_4 = 1, \Delta_5 = 0.285, \Delta_6 = -2, \Delta_7 = 0.001$. Singular periodic solutions of Eq. (3.8) is introduced in Figure 2 with $c = -2.95, \Delta_1 = -0.565, \Delta_3 = -0.025, \Delta_4 = -0.615, \Delta_5 = 0.04, \Delta_6 = 0.46, \Delta_7 = -0.02$. Figure 3 shows a singular solitary solution of Eq. (3.11) with $c = 0.02, \Delta_1 = -2, \Delta_2 = -2, \Delta_3 = -2, g_2 = -2$. Figure 4 shows a dark solitary solution of Eq. (3.16) with $c = -0.48, a_1 = -0.22, g_4 = 0.02, g_3 = -0.26, \Delta_3 = -0.15, \Delta_4 = 0.86, \Delta_5 = 0.1, \Delta_7 = -0.28$.

5. DISCUSSION AND RESULTS

This paper successfully ventured and recovered optical soliton solutions to the dispersive concatenation model with linear chromatic dispersion and self-phase modulation. Earlier investigations have laid the groundwork by introducing concatenation models combining established equations such as the NLSE, LPDE model, and Sasa-Satsuma equation. We have extended these models to incorporate higher-order dispersive effects, introducing equations like the SHE and quintic-order NLSE. Studying was conducted with the aid of the improved modified extended tanh function method, various and novel solutions were raised. These solutions including bright, dark and singular solitons. In addition, other mathematical solutions such as Weierstrass elliptic, exponential, rational and singular periodic solutions were constructed. 3D and 2D graphical representations were illustrated for some selected solutions to show the nature of the propagated waves. These solutions were presented by setting the parameters with appropriate values. The obtained






FIGURE 1. Bright solitary solution of Eq. (3.7).




FIGURE 2. Singular periodic solution of Eq. (3.8).

solitons proved that a dedicated balance occurred between the non linear and dispersion terms. These solutions will help in the development of the communication industry as these types of solutions can propagate to very long distances maintaining their shapes and speeds.

6. CONCLUSION

The IMETS was successfully applied in this work to investigate dispersive optical solitons. Many solitons and other solutions were extracted. These solutions including bright, dark, and singular solitary solutions, Weierstrass elliptic and singular periodic solutions. Moreover, graphical representations in both 2D and 3D of some of the recovered solutions are presented to illustrate the characteristics of the propagating wave. These solutions provide an explanation for a wide range of fascinating and challenging physical phenomena due to the NLSE model's applicability in several scientific domains, including wave-guides and optical fibers. The retrieved solutions in this research study are novel, and the model was not previously investigated using the proposed methodology. The approach's success, convenience of use, and efficacy show the method's applicability for dealing with nonlinear optical problems. With all of these




FIGURE 3. Singular solitary solution of Eq. (3.11).




FIGURE 4. Dark solitary solution of Eq. (3.16).

features, it will undoubtedly enrich the literature. The results are thus tremendously promising and lead to the avenues of further research in this arena. Later, the model will be studied with differential group delay followed by the consideration of the model with dispersion-flattened fibers. In addition, the stochastic model can be investigated to show the effect of the noise on the extracted solutions.

REFERENCES

- [1] I. Samir, N. Badra, A. R. Seadawy, H. M. Ahmed, A. H. Arnous, *Computational extracting solutions for the perturbed Gerdjikov-Ivanov equation by using improved modified extended analytical approach*, Journal of Geometry and Physics, 176 (2022), 104514.
- [2] I. Onder, A. Secer, M. Ozisik, and M. Bayram, *Investigation of optical soliton solutions for the perturbed Gerdjikov-Ivanov equation with full-nonlinearity*, Heliyon, 9 (2) (2023), e13519.
- [3] S. Arshed, A. Biswas, P. Guggilla, and A. S. Alshomrani, *Optical solitons for Radhakrishnan–Kundu–Lakshmanan equation with full nonlinearity*, Physics Letters A, 384(26) (2020), 126191.
- [4] O. González-Gaxiola and A. Biswas, *Optical solitons with Radhakrishnan–Kundu–Lakshmanan equation by Laplace–Adomian decomposition method*, Optik, 179 (2019), 434–42.
- [5] N. A. Kudryashov, *On traveling wave solutions of the Kundu–Eckhaus equation*, Optik, 224 (2020), 165500.



- [6] S. Chen and Z. Yan, *The Hirota equation: Darboux transform of the Riemann–Hilbert problem and higher-order rogue waves*, Applied Mathematics Letters, 95 (2019), 65–71.
- [7] H. H. Hussein, H. M. Ahmed, and W. Alexan, *Analytical soliton solutions for cubic-quartic perturbations of the Lakshmanan–Porsezian–Daniel equation using the modified extended tanh function method*, Ain Shams Engineering Journal, 15 (3) (2024), 102513.
- [8] A. Ankiewicz and N. Akhmediev, *Higher-order integrable evolution equation and its soliton solutions*, Physics Letters, 378 (4) (2014), 358–361.
- [9] A. Ankiewicz, Y. Wang, S. Wabnitz, and N. Akhmediev, *Extended nonlinear Schrödinger equation with higher-order odd and even terms and its rogue wave solutions*, Physical Review E, 89(1) (2014), 012907.
- [10] A. Chowdury, D. Kedziora, A. Ankiewicz, and N. Akhmediev, *Breather-to-soliton conversions described by the quintic equation of the nonlinear Schrödinger hierarchy*, Physical Review E, 91(3) (2015), 032928.
- [11] A. Chowdury, D. Kedziora, A. Ankiewicz, and N. Akhmediev, *Soliton solutions of an integrable nonlinear Schrödinger equation with quintic terms*, Physical Review E, 90(3) (2014), 032922.
- [12] A. Chowdury, D. Kedziora, A. Ankiewicz, and N. Akhmediev, *Breather solutions of the integrable quintic nonlinear Schrödinger equation and their interactions*, Physical Review E, 91(2) (2015), 022919.
- [13] W. B. Rabie, H. M. Ahmed, A. R. Seadawy, A. Althobaiti, *The higher-order nonlinear Schrödinger’s dynamical equation with fourth-order dispersion and cubic-quintic nonlinearity via dispersive analytical soliton wave solutions*, Optical and Quantum Electronics, 53 (2021), 1–25.
- [14] E. M. Zayed, K. A. Gepreel, M. El-Horbaty, A. Biswas, Y. Yıldırım, H. Triki, and A. Asiri, *Optical Solitons for the Dispersive Concatenation Model*, Contemporary Mathematics, 4(3) (2023), 592–611.
- [15] I. Samir, H. M. Ahmed, M. Mirzazadeh, and H. Triki, *Derivation new solitons and other solutions for higher order Sasa–Satsuma equation by using the improved modified extended tanh scheme*, Optik, 274 (2023), 170592.
- [16] O. El-shamy, R. El-barkoki, H. M. Ahmed, W. Abbas, and I. Samir, *Exploration of new solitons in optical medium with higher-order dispersive and nonlinear effects via improved modified extended tanh function method*, Alexandria Engineering Journal, 68 (2023), 611–618.
- [17] A. R. Seadawy, *Stability analysis for Zakharov–Kuznetsov equation of weakly nonlinear ion-acoustic waves in a plasma*, Computers & Mathematics with Applications, 67(1) (2014), 172–180.
- [18] A. R. Seadawy, K. K. Ali, and R. Nuruddeen, *A variety of soliton solutions for the fractional Wazwaz–Benjamin–Bona–Mahony equations*, Results in Physics, 12 (2019), 2234–2241.
- [19] A. R. Seadawy, S. T. Rizvi, I. Ali, M. Younis, K. Ali, M. Makhlof, and A. Althobaiti, *Conservation laws, optical molecules, modulation instability and Painlevé analysis for the Chen–Lee–Liu model*, Optical and Quantum electronics, 53 (2021), 1–15.
- [20] A. R. Seadawy, M. Arshad, and D. Lu, *The weakly nonlinear wave propagation theory for the Kelvin–Helmholtz instability in magnetohydrodynamics flows*, Chaos, Solitons & Fractals, 139 (2020), 110141.
- [21] A. Jhangeer, H. Rezazadeh, and A. Seadawy, *A study of travelling, periodic, quasiperiodic and chaotic structures of perturbed Fokas–Lenells model*, Pramana, 95 (2021), 1–11.
- [22] A. R. Seadawy and B. A. Alsaedi, *Soliton solutions of nonlinear Schrödinger dynamical equation with exotic law nonlinearity by variational principle method*, Optical and Quantum Electronics, 56(4) (2024), 700.
- [23] A. R. Seadawy and B. Alsaedi, *Contraction of variational principle and optical soliton solutions for two models of nonlinear Schrodinger equation with polynomial law nonlinearity*, AIMS Math, 9(3) (2024), 6336–6367.

