

## Phenotypic and genotypic characterization of extended-spectrum $\beta$ -lactamase genes in *Pseudomonas aeruginosa* isolates from local dogs bred in Nsukka, Nigeria

Adonu Cyril Chekwue<sup>1</sup>, Ezema James Nnabuike<sup>2\*</sup>, Ujam Treasure Nonye<sup>1</sup>, Ali Ibeabuchi<sup>3</sup>, Okorie Nndiamaka Hannah<sup>3</sup>, Onwusoba Restus Chibuiké<sup>4</sup>, Onyi Patrick<sup>4</sup>

<sup>1</sup>Department of Pharmaceutical Microbiology and Biotechnology, Faculty of Pharmaceutical Sciences, Enugu State University of Science and Technology, Enugu State, Nigeria

<sup>2</sup>Department of Medical Microbiology, Faculty of Basic Clinical Sciences, Enugu State University of Science and Technology, Enugu State, Nigeria

<sup>3</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Enugu State University of Science and Technology, Enugu State, Nigeria

<sup>4</sup>Department of Pharmaceutical Microbiology and Biotechnology, Faculty of Pharmaceutical Sciences, University of Nigeria Nsukka, Enugu State, Nigeria

### Article type:

Original article

### Keywords:

*Pseudomonas aeruginosa*

Dogs

PCR

Dog

ESBL genes

Nsukka

### Article history:

Received:

July 23, 2024

Revised:

November 16, 2024

Accepted:

February 10, 2025

Available online:

February 13, 2025

### Abstract

One of the major public health concerns is the transmission of antibiotic-resistant bacteria including extended spectrum beta-lactamase (ESBL)-producing *Pseudomonas aeruginosa* (*P. aeruginosa*) from animals to humans. This study investigated the prevalence of blaSHV, blaCTX-M, blaOXA and blaTEM ESBL genes in *P. aeruginosa* isolates from Nigeria local dogs bred in Nsukka. Anal swab samples (n = 150) were bacteriologically analyzed on soybean casein digest broth and cetrimide agar for the enrichment and isolation of *P. aeruginosa* respectively. Antibiotic sensitivity test was done according to Clinical and Laboratory Standard Institute (CLSI) criteria. Extended-Spectrum B-Lactamases production was detected phenotypically using the double disc synergy test (DDST) method and genotypically multiplex PCR technique respectively. A total of 39 (26.0 %) of the 150 samples were *P. aeruginosa* which were multiply resistant to antibiotics tested. The *P. aeruginosa* isolates were found to be multiply resistant to antibiotics tested. Twenty-One (53.9%) of the isolates showed ESBL phenotype while 18 (46.2 %) of the isolates were confirmed ESBL positive by Polymerase Chain Reaction (PCR). The prevalence of blaSHV, blaCTX-M, blaOXA and blaTEM genes in *P. aeruginosa* tested were 4 (10.3%), 17 (43.6 %), 15 (38.5%), and 6 (15.4%), respectively. This study, therefore, reported multi-drug resistant ESBL-positive *P. aeruginosa* in these local dog breeds. The exposure of these animals to antibiotics accumulates antibiotic resistance determinants in their intestinal flora.

\*Corresponding author: [nnabuike.ezema@esut.edu.ng](mailto:nnabuike.ezema@esut.edu.ng)

<https://doi.org/10.22034/jzd.2024.62579.1275>

[https://jzd.tabrizu.ac.ir/article\\_19286.html](https://jzd.tabrizu.ac.ir/article_19286.html)

Cite this article: Chekwue AC, Nnabuike EJ, Nonye UT, Ibeabuchi A, Hannah ON, Chibuiké OR, Patrick O. Phenotypic and genotypic characterization of extended-spectrum  $\beta$ -lactamase genes in *Pseudomonas aeruginosa* isolates from local dogs bred in Nsukka, Nigeria. Journal of Zoonotic Diseases. 2025, 9 (2): 781-791.

Copyright© 2025, Published by the University of Tabriz.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY NC)



## Introduction

*Pseudomonas aeruginosa* (*P. aeruginosa*) is an opportunistic Gram-negative pathogen (1), colonizing diverse environments owing to its ability to dwell on a variety of organic food sources. The bacterium causes infection in some animals such as sheep, horses, bovines and dogs (2). In humans, it causes dermatitis, meningitis, skin infections in severe burns, sepsis, and nosocomial infections of the urinary tract (3, 4). This bacterium is also implicated in eye and digestive system infections (5). *P. aeruginosa* is reported as a common pathogen of both community (non-hospital) and hospital environments with the ability to form biofilms and resist antibiotics (6, 7). Thus, the bacterium is multidrug-resistant (MDR) in nature (8). Recently, the prevalence of MDR *P. aeruginosa* strains has been on the increase, with rates between 15% and 30% (9, 10). One of the major upshots of multidrug resistance is the difficulty of selecting an appropriate empirical antibiotic treatment. Patients with infections caused by MDR *P. aeruginosa* are prone to inadequate antibiotic treatment (11). When there is delay in receiving the right antibiotic therapy, the resultant effects include very high morbidity and mortality rates in patients with *P. aeruginosa* infections (12). The most common mechanism of resistance among the Gram-negative pathogens is the production of hydrolytic enzymes, the beta-lactamases. (13). Gram-negative bacteria that produce beta-lactamases are a major concern in healthcare due to their ability to spread globally. Complications in many bacterial infections are on the increase because of the increasing prevalence of beta-lactamases producing pathogens (14). The case is even worse with a class of enzymes called extended-spectrum beta-lactamases (ESBLs) which are enzymes that destroy most beta-lactam antibiotics, including cephalosporins and monobactams (15). The commonly encountered groups of these ESBL enzymes are TEM, SHV, OXA and CTX-M (15). The production of these enzymes is primarily plasmid encoded with multiple resistance genes including those of non-

beta-lactam antibiotics (16-18), leaving clinicians with a limited choice of drugs in both human and veterinary medicine. Compared with human medicine, there is a paucity of information concerning ESBL-producing bacteria in veterinary medicine. The activity of these ESBL enzymes is inhibited by clavulanic acid, sulbactam, and tazobactam, and this phenomenon is utilized in the phenotypic detection of ESBLs. ESBL enzyme production is known in human clinical cases (19). Consequently, animals that are in close contact with humans contract ESBL-positive microorganisms in a reverse zoonotic event, called zoonoanthroposis (20). Thus, food and companion animals are frequently implicated as potential reservoirs of multi-antibiotic-resistant pathogens (21). In Nsukka, Nigeria, local dogs serve as both companion and food animals. On a daily basis, these animals are slaughtered for business purposes just like any other farm animal. Their meats are consumed by the indigenes. So, humans are always in contact with the local dogs as either their companions or animals reared for their meat. The knowledge of the prevalence of *P. aeruginosa* in local dogs is important owing to its zoonotic and gene transfer potentials. The spread of these "high-risk" clones poses a threat to global public health that needs urgency and determination (21), thus, justifying the need for its study in the non-hospital environment to prevent any hazard that may be caused by them. In addition to monitoring the antimicrobial resistance patterns of *P. aeruginosa* isolates from both humans and animals, it is pertinent to elucidate the antibiotic resistance mechanism of these organisms to preserve global public health. Currently, there is no data on the genotypic characteristics of ESBL genes in *P. aeruginosa* from Nigerian local dogs in Nsukka. This is because adequate attention has not been given to the studies involving ESBL in *P. aeruginosa* isolates from local dogs in the area, possibly due to fear of the aggressive nature of some of these animals. Therefore, the objective of this study was to characterize, by phenotypic and

genotypic methods, the ESBL genes among *P. aeruginosa* from Nigerian local dogs bred in Nsukka, South-East, Nigeria.

### Materials and methods

#### *Isolation and characterization of P. aeruginosa*

One hundred and fifty (150) anal swab samples were used for the study and all the samples were analyzed using standard microbiology procedures. The samples were inoculated in soybean casein digest broth (Oxoid, UK) and incubated for 24 h at 37°C. Bacterial growth was suspected due to the turbidity observed after incubation. Each colony was sub-cultured on ceftrimide selective agar (Oxoid, UK) for the selective isolation of *P. aeruginosa* (22). Suspension of broth culture showing turbidity was aseptically streaked onto ceftrimide agar plates for the selective isolation of *P. aeruginosa*. The plates were incubated for 24 – 48 h at 37°C. Suspected *P. aeruginosa* isolates were further purified by culturing on ceftrimide agar, the presence of which was confirmed by characteristic colonial morphology, Gram staining, growth at 42 °C / 48 h in soybean casein digest broth and biochemical testing (23).

#### *Phenotypic detection of the ESBL enzymes and antibiotic sensitivity testing*

Inhibition diameter of less than 26 mm for both cefotaxime and ceftriaxone were selected for the Double Disc Synergy Test (DDST) for ESBL production by using a disc of amoxicillin-clavulanate (20/10 µg) along with two other antibiotics belonging to the third generation cephalosporins (3GC); cefotaxime (30 µg) and ceftriaxone (30 µg) (24) with some modifications. A liquid culture of each of all the bacteria to be analyzed (*P. aeruginosa* strains and the controls) was prepared by adjusting the prepared bacterial suspension to 0.5 McFarland opacity standard. Then, a lawn culture of the organisms was prepared by pouring the standardized liquid culture on a dried Mueller-Hinton agar plate and decanting the excess from the plate. An antibiotic disc of amoxicillin-clavulanic acid (20/10 µg) was placed at the center

of the MH agar plate, and antibiotic discs containing cefotaxime (30 µg) and ceftazidime (30 µg) were each placed at a distance of 15mm (center to center) from the central disc, amoxicillin-clavulanic acid 20/10 µg. The plates were incubated at 37°C for 18–24 h. Phenotypic inference for ESBL production was drawn from the expansion of the zones of inhibition of the cephalosporins by the amoxicillin-clavulanic acid disc (20/10 µg). A ≥5mm increase in the inhibition zone diameter for either of the cephalosporins (ceftazidime and cefotaxime) tested in combination with amoxicillin-clavulanic acid versus its zone when tested alone confirms ESBL production phenotypically. The positive and negative controls for ESBL production were *Klebsiella pneumoniae* 700603 and *Escherichia coli* 25922 respectively. The *P. aeruginosa* isolates were tested for their antibiotic susceptibilities by the disc diffusion method according to the Clinical and Laboratory Standard Institute (CLSI) guidelines (25). The following antibiotics were used; amoxicillin-clavulanic acid (20/10 µg), meropenem (10 µg), cefotaxime (30 µg), ceftazidime (30 µg), sulfamethoxazole-trimethoprim (25 µg), gentamicin (10 µg), ceftriaxone (30 µg), ciprofloxacin (10 µg), ofloxacin (10 µg), ampicillin (10 µg) and cloxacillin (10 µg). Other classes of antibiotics were used to test for multidrug resistance. All the antibiotic discs were sourced from Oxoid, UK. The diameter of the inhibition zone was measured and interpreted based on the CLSI criteria.

#### *Screening for ESBL gene using PCR method*

The phenotypically confirmed ESBL positive *P. aeruginosa* strains stored in slants were sub-cultured on nutrient agar plates and then on nutrient broth (Oxoid, UK) prior to DNA extraction. The pure single colony of the isolates was further subcultured in 5 ml of nutrient broth and incubated at 37°C overnight for DNA extraction. Genomic DNA of the test isolate was prepared using ZR fungal/bacterial DNA miniprep (Zymo Research, California) following manufacturer's instructions. Primers used for amplification of resistance genes

are shown in Table 1. ESBL-positive *P. aeruginosa* was examined for the presence of TEM, SHV, OXA-1 and CTX-M-1 genes by PCR technique. ESBL genes were amplified using a thermal cycler (Lumex instruments, Canada) with a final volume of 26.5 µL master mix comprising 0.2 µL of Taq polymerase enzyme U/µL, 2.5 µL of 10X PCR buffer along with 2.5 µL MgCl<sub>2</sub>, 1 µL of 10 pM from each of the forward and reverse primers, 2.5 µL of dNTPs MIX (2 mM), 3 µL of DNA template (from the test isolates), 14.8 µL of nuclease-free water. Cycling conditions include initial denaturation at 94°C for 5 min, 36 cycles of denaturation at 94°C for 30 s and an initial

elongation step at 72°C for 45 s, followed by the final elongation step at 72°C for 7 min.

#### *Gel electrophoresis*

Amplified PCR products were separated on 1.5 % agarose gel (3 g of the agarose powder/ 0.5X TBE buffers), according to standard procedure (24). The gel was run at 100 V for 1.5 h; and then visualized with UV transilluminator at 260 nm.

#### *Statistical analysis*

Data analysis was carried out using the Statistical Package for Social Sciences (SPSS) version 23.0 (SPSS, Chicago, IL, USA). Data were analyzed in terms of percentages.

**Table 1.** Primer sequence for ESBL gene characterization

Gene	Primer sequence	Expected amplicon size (bp)
<i>bla</i> TEM F	5'AAACGCTGGTGAAAGTA3'	500
<i>bla</i> TEM R	5'AGTGTGTTTAGAATGGTGATC3'	
<i>bla</i> OXA F	5'ACACAATACATATCAACTTCGC3'	650
<i>bla</i> OXA R	5'AGTGTGTTTAGAATGGTGATC3'	
<i>bla</i> CTX-M F	5'CGCTTTGCGATGTGCAG3'	750
<i>bla</i> CTX-M R	5'ACCGCGATATCGTTGGT3'	
<i>bla</i> SHV F	5'ATGCGTTATATTCGCCTGTG3'	850
<i>bla</i> SHV R	5'TGCTTTGTTATTCGGGCCAA3'	

## Results

### *Prevalence of P. aeruginosa and ESBL genes*

A total of 39 (26.0%) *P. aeruginosa* isolates were recovered from 150 anal swab samples used in the study. Isolates showed characteristic blue-green pigments which diffused into the medium, positive reaction to Oxidase test and growth at 42 °C / 48 h in soybean casein digest broth. The presence of ESBL phenotypes in 21 (53.9%) out of the 39 isolates shows the appearance of ESBL positive and negative bacteria when cultivated on Mueller Hinton agar (MHA) in the presence of third

generation cephalosporins (cefotaxime and ceftriaxone) and amoxicillin-clavulanic acid (Table 2).

### *Antibiotic susceptibility pattern of P. aeruginosa isolates and the prevalence of the encoding ESBL genes*


The *P. aeruginosa* isolates exhibited different levels of sensitivity to the test antibiotics. The susceptibility patterns, as well as the ESBL genes encoding the resistance, are shown in Table 3.

**Table 2.** Prevalence of ESBL genes in the 39 isolates of *P. aeruginosa* from anal swab samples


Number of isolates n (%)	ESBL gene harboured
17(43.6 %)	<i>Bla</i> CTX-M
15(38.5%)	<i>Bla</i> OXA
6(15.4 %)	<i>Bla</i> TEM
4(10.3%)	<i>Bla</i> SHV

**Table 3.** Antibiotic susceptibility pattern of *P. aeruginosa* isolates and the distribution of the encoding ESBL genes (n = 39)


Antibiotics	Susceptibility pattern			Encoding ESBL gene
	Resistant	n (%) Intermediate	Sensitive	
Amoxicillin-clavulanic acid	22 (56.4)	8 (20.5)	9 (23.1)	<i>Bla</i> CTX-M, <i>bla</i> TEM, <i>bla</i> SHV
Meropenem	6 (15.3)	3(7.7)	30(77.0)	<i>Bla</i> OXA
Cefotaxime	20(51.3)	15(38.5)	4 (10.3)	<i>Bla</i> CTX-M, <i>bla</i> OXA
Ceftazidime	30 (77.0)	6 (15.3)	3(7.7)	<i>Bla</i> CTX-M, <i>bla</i> OXA
Sulfamethoxazole-Trimethoprim	27(69.2)	11(28.2)	1 (2.6)	<i>Bla</i> CTX-M, <i>bla</i> OXA, <i>bla</i> TEM, <i>bla</i> SHV
Gentamicin	20(51.3)	5(12.8)	14(35.9)	<i>Bla</i> CTX M, <i>bla</i> OXA, <i>bla</i> TEM, <i>bla</i> SHV
Ceftriaxone	20(51.3)	1(2.6)	18(46.0)	<i>Bla</i> CTX-M, <i>bla</i> OXA
Ciprofloxacin	20(51.3)	9(23.1)	10(25.6)	<i>Bla</i> CTX-M, <i>bla</i> TEM, <i>bla</i> SHV
Ofloxacin	20(51.3)	10(25.6)	9(23.1)	<i>Bla</i> CTX-M, <i>bla</i> OXA, <i>bla</i> TEM, <i>bla</i> SHV
Ampicillin	39 (100)	0 (0)	0 (0)	<i>Bla</i> TEM, <i>bla</i> SHV
Cloxacillin	38 (97.4)	1 (2.6)	0 (0)	<i>Bla</i> CTX-M, <i>bla</i> OXA, <i>bla</i> TEM, <i>bla</i> SHV




**Fig. 1.** Gel image of the amplified *bla* CTX-M gene showing amplification at 750bp. M is a 50bp ladder used for estimation of the size band of amplicons.



**Fig. 2.** Gel image of *bla* OXA gene showing amplification at about 650bp. M is a 50bp ladder used for estimation of the size band of amplicons.



**Fig. 3.** Gel image of *bla* TEM gene showing amplification of TEM at about 500bp. M is a 50bp ladder used for estimation of the size band of amplicons.



**Fig. 4.** Gel image of *bla* SHV gene showing amplification of SHV at about 850bp. M is a 50bp ladder used for estimation of the size band of amplicons.

### Discussion

Antibiotics are important agents for the management of diseases caused by bacteria in both human and veterinary medicine (26). However, the increased emergence of antibiotic-resistant bacteria, including those that produce ESBLs have rendered many of these agents ineffective (13, 15). As a result, many of the multi-antibiotic resistant organisms remain viable even in the face of broad spectrum antibiotics (15). The need to search for the possible reservoirs of resistant organisms including ESBL producers from the non-hospital milieu ignited the zeal for this study. We, therefore, investigated the prevalence of ESBL phenotypes and genotypes; particularly *bla*SHV, *bla*CTX-M, *bla*OXA and *bla*-TEM genes in *P. aeruginosa* isolates from Nigeria local dogs bred in Nsukka. The isolation rate of *P. aeruginosa* in this study was high ( $n = 39$ , 26.0%), considering the nature and severity of diseases associated with this organism (Table 2). This result is not in agreement with the results of the work done by other researchers who isolated 16.7% and 17.7 % of *P. aeruginosa* strains from samples collected from healthy and sick dogs respectively (12). Similarly, several other reports of

research carried out on dogs and other related animals revealed the presence of *P. aeruginosa* in these animals (27-29). In a related study conducted in Egypt (28), the prevalence of *P. aeruginosa* isolates in camel meat was about 22.5%, confirming the presence of this organism in other animals. With these results of the isolation rates of *P. aeruginosa* in samples from dogs, understanding the connections between *P. aeruginosa* and dogs is, therefore, very important for many reasons. Apart from helping veterinary doctors to choose the most effective antibiotic options for the treatment of sick dogs, understanding the transmission and prevention of *P. aeruginosa* infections in dogs can have implications for public health, especially as dogs can act as reservoirs of *P. aeruginosa*. In the present study, the antibiotic sensitivity test result of ESBL producing *P. aeruginosa* showed that these isolates were highly resistant to beta-lactam antibiotics in the penicillin family including ampicillin (100%), amoxicillin-clavulanic acid (56.4%) and cloxacillin (97.4%; Table 3) This is in agreement with the results of the work conducted in South Africa, which showed that most *P. aeruginosa* isolates from dogs were resistant to a wide range of antibiotics (30). A high proportion of

MDR *P. aeruginosa* isolates was resistant to Amoxicillin-Clavulanic acid (99%). This pattern was followed by resistance to antibiotics in the Cephalosporin family including Cefotaxime (51.3%), Ceftazidime (77.0%) and Ceftriaxone (51.3%). The reduced susceptibility of the isolates to the Cephalosporins as observed in the present work is similar to the results of the study conducted by (31, 29) in southwest Nigeria and Abakaliki respectively. Furthermore, Carbapenem class of beta-lactams represented by Meropenem showed 15.3 % resistance. A similar result showed that a low proportion of *P. aeruginosa* isolates were resistant to Imipenem (6%) (30). For non-beta lactam antibiotics, increased resistance of the isolates to fluoroquinolones (Ciprofloxacin and Ofloxacin) (51.3 %) and Gentamicin (51.3%) was recorded in this study, which is in agreement with the results of the same work conducted in South Africa (30). Conversely, a dissimilar result on the fluoroquinolone, cephalosporin and Gentamicin resistance profile of *P. aeruginosa* isolates from dogs was given in Japan because their findings showed lower resistance rates of Ciprofloxacin, Cefotaxime and Gentamicin which were 20.5%, 17.8% and 4.1%, respectively (32). Furthermore, such lower resistant rates to aminoglycosides and fluoroquinolones such as Ciprofloxacin (40.5%) have been reported in Malaysia (33) and India (34) respectively. *Pseudomonas aeruginosa* is remarkably resistant to antibiotics due to the outer membrane-associated permeability barrier of the organism (35). It is important to note that the addition of antibiotics in animal feed to combat infections and promote growth poses a health risk to the human population due to the possible development and dissemination of antibiotic resistance (36). Also in this study, ESBL production was phenotypically detected in 53.9% of all the *P. aeruginosa* isolates. This is similar to research conducted in Italy by some researchers who detected ESBL in *P. aeruginosa* isolates from dogs (37). In a related study carried out in Egypt (28), researchers reported a lower prevalence (45%) of

phenotype of ESBL in isolates of *P. aeruginosa* from Camel. Unlike our findings, other researchers reported a much lower prevalence (20.8%) of ESBL producing clinical Enterobacteriaceae from dogs in Switzerland (38). The disparity in the prevalence of ESBL enzymes in these animals may be related to the differences in animal species, screening methods, dexterity of the researchers and the materials used for the tests. The PCR technique confirmed ESBL production in 18 (46.2%) of the *P. aeruginosa* isolates as shown in the gel images (Figures 1-4). The prevalence of blaCTX-M, blaOXA, blaTEM and blaSHV genes in *P. aeruginosa* tested were, 17 (43.6%), 15 (38.5%), 6 (15.4 %) and 4 (10.3 %), respectively. The presence of the blaCTX-M gene found in this study is in agreement with the result of the work done in Taiwan by other researchers who observed the blaCTX-M in 58.5% of the bla genes (39). Our findings on the prevalence of these ESBL genes are not in agreement with the report of the work done in India which showed that the predominant beta-lactamase genes detected in *P. aeruginosa* isolates from dogs include blaOXA (69.5%) followed by blaTEM (36.9%), bla-SHV (26.0%), blaCTX-M g (15.2%) and blaAmpC (4.34%) (40). In Egypt, it was observed that the most commonly detected  $\beta$ -lactamase-genes in *P. aeruginosa* were blaCTX-M (38%), followed by blaSHV (33.3%) and blaTEM (27). CTX-M enzymes have now been investigated to be the most disseminated type of ESBL globally and represent the wide dissemination of particular plasmids or bacterial clones (41). Both phenotypic detection and genotypic confirmation of ESBL genes in food and companion animals such as dogs is a grave challenge facing the human populace, especially in rural communities where hygiene and veterinary services are lacking.

### Conclusion

We confirmed the notoriety of *P. aeruginosa* isolates from Nigeria local dogs for the production of ESBL enzymes. The high prevalence of the enzymes was corroborated by high prevalence of blaSHV, blaCTX-M, blaOXA and blaTEM genes in

the organisms isolated from the dogs. The presence of these genes has boosted the resistance of the organism to both beta-lactams and some non-beta-lactam antibiotics. The presence of ESBL genes in the isolates detected in this study is not far from several risk factors connected with the evolution and spread of antibiotic resistance genes in the non-hospital environment. It is possible that some of these isolates recovered from dogs were contracted from environments and/or owners and could be transmitted back to the owners and vice versa. The present study reporting the first detection of ESBL resistant phenotype and genotypes in *P. aeruginosa* strains of canine origin in Nsukka, South-East Nigeria consolidates our knowledge on the alarming ESBL resistance reports in the gut microbiota of dogs world-wide. The presence and persistence of these genes in non-human sources may pose a great risk to public health. Therefore, strict surveillance of this pathogen is very necessary to assuage the emergence and spread of multidrug-resistant strains, which can pose significant challenges in clinical settings.

#### Acknowledgments

We appreciate Mr Eze Leonard and all the staff of Adonai research laboratory for their technical support. We are grateful for the assistance rendered to us by Mr Lowyaar Nwarko, Mrs Angella Lady B and others who allowed us to collect the specimen from their freshly slaughtered dogs. Their contributions are immeasurable as they greatly assisted in the completion of this work.

#### Ethical approval

The research was carried out following the World Medical Association (WMA) declaration of Helsinki on the principles for medical research involving human and animal subjects (42). There was no written ethical approval as the samples were collected at the point of slaughter.

#### Competing interests

The authors declare no competing interest.

#### References

1. Yeung AT, Torfs EC, Jamshidi F, Bains M, Wiegand I, Hancock RE, et al. Swarming of *Pseudomonas aeruginosa* is controlled by a broad spectrum of transcriptional regulators, including MetR. *J Bacteriol.* 2009;191(18):5592-602. <https://doi.org/10.1128/jb.00157-09>
2. Bricha S, Ounine K, Oulkheir S, Haloui NEEL, Attarassi B. Virulence factors and epidemiology related to *Pseudomonas aeruginosa*. *Tunisian J Infect Dis.* 2009; 2: 7–14. [https://www.infectiologie.org.tn/pdf\\_ppt\\_docs/revues/rti11/revue\\_generale1.pdf](https://www.infectiologie.org.tn/pdf_ppt_docs/revues/rti11/revue_generale1.pdf)
3. Mitov I, Strateva T, Markova B. Prevalence of virulence genes among bulgarian nosocomial and cystic fibrosis isolates of *pseudomonas aeruginosa*. *Braz J Microbiol.* 2010; 41(3): 588-95. <https://doi.org/10.1590/S1517-83822010000300008>.
4. Rostamzadeh Z, Mohammadian M, Rostamzade A. Investigation of *Pseudomonas aeruginosa* resistance pattern against antibiotics in clinical samples from an Iranian educational hospital. *Adv Microbiol.* 2016; 6: 190-4. <https://doi.org/10.4236/aim.2016.63019>.
5. Gellatly SL, Hancock RE. *Pseudomonas aeruginosa*: new insights into pathogenesis and host defenses. *Pathog Dis.* 2013; 67(3): 159-73. <https://doi.org/10.1111/2049632X.12033>
6. Brooks GF, Butel JS, Morse SA. *Medical Microbiology*, 23rd edition. New York: McGraw Hill; 2004; 248-60.
7. Haleem JK, Tarrad T, Banyan A. Isolation of *Pseudomonas aeruginosa* from clinical cases and environmental samples, and analysis of its antibiotic resistant spectrum at Hilla Teaching Hospital. *Med J Babylon.* 2011; 8(4): 618-24.
8. Walkty A, Lagace-Wiens P, Adam H, Baxter M, Karlowsky J, Mulvey MR, et al. Antimicrobial susceptibility of 2906 *Pseudomonas aeruginosa* clinical isolates obtained from patients in Canadian hospitals over a period of 8 years: Results of the Canadian Ward surveillance study (CANWARD), 2008-2015. *Diagn Microbiol Infect Dis.* 2017; 87(1): 60-3.



- <https://doi.org/10.1016/j.diagmicrobio.2012.05.007>
9. Paterson DL, Rice LB. Empirical antibiotic choice for the seriously ill patient: are minimization of selection of resistant organisms and maximization of individual outcome mutually exclusive? *Clin Infect Dis*. 2003; 36(8): 1006-12. <https://doi.org/10.1086/374243>
  10. Sader HS, Castanheira M, Duncan LR, Flamm RK. Antimicrobial Susceptibility of Enterobacteriaceae and *Pseudomonas aeruginosa* Isolates from United States Medical Centers Stratified by Infection Type: Results from the International Network for Optimal Resistance Monitoring (INFORM) Surveillance Program, 2015-2016. *Diagn Microbiol Infect Dis*. 2018; 92(1): 69-74. <https://doi.org/10.1016/j.diagmicrobio.2018.04.012>.
  11. Guillaumet CV, Vazquez R, Noe J, Micek ST, Kollef MH. A cohort study of bacteremic pneumonia: The importance of antibiotic resistance and appropriate initial therapy? *Medicine (Baltimore)*. 2016; 95(35): e4708. <https://doi.org/10.1097/md.00000000000004708>
  12. Park Y, Oh J, Park S, Sum S, Song W, Chae J, et al. Antimicrobial resistance and novel mutations detected in the *gyrA* and *parC* genes of *Pseudomonas aeruginosa* strains isolated from companion dogs. *BMC Vet Res*. 2020; 16(1): 111. <https://doi.org/10.1186/s12917-020-02328-0>.
  13. Parajuli NP, Maharjan P, Parajuli H, Joshi G, Paudel D, Sayami S, et al. High rates of multidrug resistance among uropathogenic *Escherichia coli* in children and analyses of ESBL producers from Nepal. *Antimicrob Resist Infect Control*. 2017; 11; 6: 9. <https://doi.org/10.1186/s13756-016-0168-6>.
  14. Elsayed TI, Ismail HA, Elgamal SA. The occurrence of multidrug resistant *E. coli* which produce ESBL and cause urinary tract infections. *J Appl Microbiol Biochem*. 2017; 1(1): 2-8. <https://doi.org/10.21767/2576-1412.100008>
  15. Bush K. Past and Present Perspectives on  $\beta$ -Lactamases. *Antimicrob Agents Chemother*. 2018; 62(10): e01076-18. <https://doi.org/10.1128/AAC.01076-18>.
  16. Leinberger D, Grimm V, Rubtsova M, Weile J, Schröppel K, Wichelhaus T. et al. Integrated detection of extended-spectrum-beta-lactam resistance by DNA microarray-based genotyping of TEM, SHV, and CTX-M genes. *J Clin Microbiol*. 2010; 48(2): 460-71. <https://doi.org/10.1128/JCM.00765-09>.
  17. Enwuru CA, Iwalokun B, Enwuru NV, Ezechi O, Idika N, Oluwadun A. The occurrence and modified method for phenotypic identification of ambler group A and B extended spectrum  $\beta$ -lactamases production in urino-genital gram-negative bacterial isolates from, Nigeria. *Nat Sci*. 2013; 11 (8), 1-6.
  18. Alyamani EJ, Khiyami AM, Booq RY, Majrashi MA, Bahwerth FS, Rechkina E. The occurrence of ESBL-producing *Escherichia coli* carrying aminoglycoside resistance genes in urinary tract infections in Saudi Arabia. *Ann Clin Microbiol Antimicrob*. 2017; 16(1): 1-13 <https://doi.org/10.1186/s12941-016-0177-6>.
  19. Bradford PA. Extended-spectrum beta-lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin Microbiol Rev*. 2001; 14(4): 933-51. <https://doi.org/10.1128/CMR.14.4.933-951>.
  20. Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Herman L, et al. Broad-spectrum  $\beta$ -lactamases among Enterobacteriaceae of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol Rev*. 2010 May; 34 (3): 295-316. <https://doi.org/10.1111/j.15746976.2009.00198.x>
  21. Hafsat AG, Yaqub AG, Abubakar S, Isa AG, Roy BB. Multi-drug resistant bacteria isolated from fish and fish handlers in Maiduguri, Nigeria. *Int J Animal Vet Adv*. 2015; 7: 49-54. <https://doi.org/10.19026/ijava.7.5240>
  22. Gilligan PH. *Pseudomonas* and *Burkholderia*, p.509-519. In *Manual of Clinical Microbiology*, 6th ed. Washington, D.C. American Society for microbiology, 1995

24. Cheesbrough M. District Laboratory Practice in Tropical Countries, 2<sup>nd</sup> edition. Cambridge Uni. Press, UK. 2006; 178-187.
25. Zhang CH, Liu YL, Wang JH. Detection of ESBLs and Antimicrobial susceptibility of *Escherichia coli* isolated in Henan, China. *J Anim Vet Adv.* 2010; 9(15): 2030-4. <https://doi.org/10.3923/javaa.2010.2030.2034>
26. Barbara LZ., Clinical Laboratory Standard Institute, CLSI: Performance standards for antimicrobial disc susceptibility test. Fifteenth informational supplement, CLSI document M100-Ed14, 2004; 96.
27. Li B, Webster TJ. Bacteria antibiotic resistance: New challenges and opportunities for implant-associated orthopedic infections. *J Orthop Res.* 2018; 36(1): 22-32. <https://doi.org/10.1002/jor.23656>.
28. Schink AK, Kadlec K, Hauschild T, Brenner Michael G, et al. Susceptibility of canine and feline bacterial pathogens to pradofloxacin and comparison with other fluoroquinolones approved for companion animals. *Vet Microbiol.* 2013; 162(1): 119-26. <https://doi.org/10.1016/j.vetmic.2012.08.001>.
29. Elhariri M, Hamza D, Elhelw R, Dorgham SM. Extended-spectrum beta-lactamase-producing *Pseudomonas aeruginosa* in camels in Egypt: potential human hazard. *Ann Clin Microbiol Antimicrob.* 2017; 31: 16 -21. <https://doi.org/10.1186/s12941-017-0197>.
30. Ejikegwu C, Esimone C, Iroha, I, Eze P, Ugwu M, Adikwu, M. Genotypic and phenotypic characterization of MBL genes in *Pseudomonas aeruginosa* isolates from the non-hospital environment. *J Pure & Appl Microbiol.* 2018; 12(4): 1877-85. <http://doi.org/10.22207/JPAM.12.4.23>
31. Eliasi UL, Sebola D, Oguttu JW, Qekwana DN. Antimicrobial resistance patterns of *Pseudomonas aeruginosa* isolated from canine clinical cases at a veterinary academic hospital in South Africa. *J S Afr Vet Assoc.* 2020; 91(0):e1-e6. <https://doi.org/10.4102/jsava.v91i0.2052>.
32. Aibinu I, Nwanneka T, Odugbemi T. Occurrence of ESBLs and MBL in clinical isolates of *Pseudomonas aeruginosa* from Lagos, Nigeria. *J Amer Sci.* 2007; 3(4): 81-85. <https://api.semanticscholar.org/CorpusID:212483990>
33. Harada K, Arima S, Niina A, Kataoka Y, Takahashi T. Characterization of *Pseudomonas aeruginosa* isolates from dogs and cats in Japan: current status of antimicrobial resistance and prevailing resistance mechanisms. *Microbiol Immunol.* 2012; 56(2): 123-7. <https://doi.org/10.1111/j.1348-0421.2011.00416.x>.
34. Fazlul MK, Zaini MZ, Rashid MA, Nazmul MH. Antibiotic susceptibility profiles of clinical isolates of *Pseudomonas aeruginosa* from Selayang Hospital, Malaysia. *J Biomed Res.* 2011; 22(3): 263.
35. Ramana BV, Chaudhury A. Antibiotic resistance pattern of *Pseudomonas aeruginosa* isolated from healthcare associated infections at a tertiary care hospital. *J Scient Soc.* 2012; 39(2): 78. <https://doi.org/10.4103/0974-5009.101850>.
36. Fernández M, Conde S, de la Torre J, Molina-Santiago C, Ramos JL, Duque E. Mechanisms of resistance to chloramphenicol in *Pseudomonas putida* KT2440. *Antimicrob Agents Chemother.* 2012; 56(2):1001-9. <https://doi.org/10.1128/AAC.05398-11>.
37. Shea KM; American Academy of Pediatrics Committee on Environmental Health; American Academy of Pediatrics Committee on Infectious Diseases. Nontherapeutic use of antimicrobial agents in animal agriculture: implications for pediatrics. *Pediatrics.* 2004; 114(3): 862-8. <https://doi.org/10.1542/peds.2004-1233>.
38. Gentilini F, Turba ME, Pasquali F, Mion D, Romagnoli N, Zambon E, et al. Hospitalized Pets as a Source of Carbapenem-Resistance. *Front Microbiol.* 2018; 6(9): 2872. <https://doi.org/10.3389/fmicb.2018.0287>
39. Zogg AL, Simmen S, Zurfluh K, Stephan R, Schmitt SN, Nüesch-Inderbinen M. High

Prevalence of Extended-Spectrum  $\beta$ -Lactamase Producing Enterobacteriaceae Among Clinical Isolates from Cats and Dogs Admitted to a Veterinary Hospital in Switzerland. *Front Vet Sci.* 2018; 27; 5: 62. <https://doi.org/10.3389/fvets.2018.00062>.

40. Huang YH, Kuan NL, Yeh KS. Characteristics of Extended-Spectrum  $\beta$ -Lactamase-Producing *Escherichia coli* From Dogs and Cats Admitted to a Veterinary Teaching Hospital in Taipei, Taiwan from 2014 to 2017. *Front Vet Sci.* 2020; 6(7): 395. <https://doi.org/10.3389/fvets.2020.00395>.
  41. Mohammad RN, Sreedevi B, Chaitanya RK, Srilatha CH. Detection of extended spectrum beta-lactam (ESBL) resistance in *Pseudomonas* species of canine origin. *Pharma Inno.* 2017; 6(9): 89-93. <https://doi.org/10.22271/tpi>.
  42. Cantón R, Novais A, Valverde A, Machado E, Peixe L, Baquero F, et al. Prevalence and spread of extended-spectrum beta-lactamase-producing Enterobacteriaceae in Europe. *Clin Microbiol Infect.* 2008; 14 Suppl 1: 144-53. <https://doi.org/10.1111/j.1469-0691.2007.01850.x>.
  43. World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA.* 2013; 310(20): 2191-4. <https://doi.org/10.1001/jama.2013.281053>.
-